

# TIMELIKE SURFACES OF CONSTANT MEAN CURVATURE $\pm 1$ IN ANTI-DE SITTER 3-SPACE $\mathbb{H}_1^3(-1)$

SUNGWOOK LEE

*This paper is dedicated to my sensei professor Kinetsu Abe.*

ABSTRACT. It is shown that timelike surfaces of constant mean curvature  $\pm 1$  in anti-de Sitter 3-space  $\mathbb{H}_1^3(-1)$  can be constructed from a pair of Lorentz holomorphic and Lorentz antiholomorphic null curves in  $\mathbb{PSL}_2\mathbb{R}$  via Bryant type representation formulae. These Bryant type representation formulae are used to investigate an explicit one-to-one correspondence, the so-called *Lawson-Guichard correspondence*, between timelike surfaces of constant mean curvature  $\pm 1$  in  $\mathbb{H}_1^3(-1)$  and timelike minimal surfaces in Minkowski 3-space  $\mathbb{E}_1^3$ . The hyperbolic Gauß map of timelike surfaces in  $\mathbb{H}_1^3(-1)$ , which is a close analogue of the classical Gauß map is considered. It is discussed that the hyperbolic Gauß map plays an important role in the study of timelike surfaces of constant mean curvature  $\pm 1$  in  $\mathbb{H}_1^3(-1)$ . In particular, the relationship between the Lorentz holomorphicity of the hyperbolic Gauß map and timelike surface of constant mean curvature  $\pm 1$  in  $\mathbb{H}_1^3(-1)$  is studied.

## 1. INTRODUCTION

It is known that surfaces of constant mean curvature  $\pm 1$  surfaces in hyperbolic 3-space  $\mathbb{H}^3(-1)$  can be constructed from holomorphic null curves in  $\mathbb{PSL}_2\mathbb{C} = \mathrm{SL}_2\mathbb{C}/\{\pm \mathrm{id}\}$  ([4], [23]), while minimal surfaces in Euclidean 3-space  $\mathbb{E}^3$  can be constructed from holomorphic null curves in  $\mathbb{C}^3$  via well-known Weierstraß-Enneper representation formula. It is also known that spacelike surfaces of constant mean curvature  $\pm 1$  in de-Sitter 3-space  $\mathbb{S}_1^3(1)$  can be constructed from holomorphic null curves in  $\mathbb{PSL}_2\mathbb{C}$  ([2], [16]), while spacelike maximal surfaces in Minkowski 3-space  $\mathbb{E}_1^3$  can be constructed from holomorphic null curves in  $\mathbb{C}^3$  via an analogue of Weierstraß-Enneper representation formula ([18], [14]). These are all related by the *Lawson-Guichard correspondence* between minimal surfaces in  $\mathbb{E}^3$  and surfaces of constant mean curvature  $\pm 1$  in  $\mathbb{H}^3(-1)$  ([15]) and the one between spacelike maximal surfaces in  $\mathbb{E}_1^3$  and spacelike surfaces of constant mean curvature  $\pm 1$  ([21]). Note that the

---

*Date:* Received July 14, 2005.

*2000 Mathematics Subject Classification.* 53A10, 53C42, 53C50.

correspondents (they are usually called the *cousins*) in different space forms satisfy the same Gauß and Mainardi-Codazzi equations.

It is interesting to see that there exists a Lawson-Guichard correspondence between timelike minimal surfaces in  $\mathbb{E}_1^3$  and timelike surfaces of constant mean curvature  $\pm 1$  in anti-de Sitter 3-space  $\mathbb{H}_1^3(-1)$ . See sections 5, 9, and 14 (appendix I) for details. In [13], J. Inoguchi and M. Toda show that timelike minimal surfaces can be constructed from a pair of Lorentz holomorphic and Lorentz antiholomorphic null curves in  $\mathbb{R}^3$  via normalized Weierstraß formula (59). Hence, one might expect a similar construction of timelike surfaces of constant mean curvature  $\pm 1$  in  $\mathbb{H}_1^3(-1)$  in terms of Lorentz holomorphic and Lorentz antiholomorphic null curves. In this paper, we prove that a pair of Lorentz holomorphic and Lorentz antiholomorphic null curves in  $\mathbb{PSL}_2\mathbb{R}$  gives rise to a timelike surface of constant mean curvature  $\pm 1$  in  $\mathbb{H}_1^3(-1)$ . Furthermore, every timelike surface of constant mean curvature  $\pm 1$  in  $\mathbb{H}_1^3(-1)$  can be constructed from a pair of Lorentz holomorphic and Lorentz antiholomorphic null curves in  $\mathbb{PSL}_2\mathbb{R}$ .

An analogue of the hyperbolic Gauß map<sup>1</sup> can be defined for timelike surfaces of constant mean curvature in  $\mathbb{H}_1^3(-1)$  and plays an important role in studying timelike surfaces of constant mean curvature  $\pm 1$  in  $\mathbb{H}_1^3(-1)$ . It is shown in section 13 that

- (1) The hyperbolic Gauß map of a Lorentz surface  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  is Lorentz holomorphic if and only if  $\varphi$  satisfies  $H = 1$  and  $Q = 0$ .
- (2) The hyperbolic Gauß map of a Lorentz surface  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  is Lorentz antiholomorphic if and only if  $\varphi$  satisfies  $H = 1$  and  $R = 0$ .

Here,  $Q$  and  $R$  are coefficients of quadratic differentials, the so-called *Hopf pairs*. They are defined in the following section.

In [3] and [22], K. Akutagawa and J. Ramanathan proved independently that

**Theorem:** *Let  $M$  be a complete spacelike surface in de Sitter 3-space  $\mathbb{S}_1^3(1)$  with constant mean curvature  $H = \pm 1$ . Then  $M$  is a totally umbilic flat surface. Moreover,  $M$  is a parabolic type surface of revolution.*

This theorem tells us that de Sitter 3-space  $\mathbb{S}_1^3(1)$  admits horosphere type spacelike surfaces. It is also interesting to see that anti-de Sitter 3-space  $\mathbb{H}_1^3(-1)$  admits horosphere type timelike surfaces. See section 10 for details.

## 2. TIMELIKE SURFACES IN ANTI-DE SITTER 3-SPACE $\mathbb{H}_1^3(-1)$

Let  $\mathbb{E}_2^4$  be the semi-Euclidean 4-space with natural coordinates  $(x_0, x_1, x_2, x_3)$  and the semi-Riemannian metric  $\langle \cdot, \cdot \rangle$  of signature  $(-, -, +, +)$  given by the quadratic form  $-(dx_0)^2 - (dx_1)^2 + (dx_2)^2 + (dx_3)^2$ .

---

<sup>1</sup>The hyperbolic Gauß map was introduced by C. Epstein in [7] and used by R. .L. Bryant to study cmc 1 surfaces in  $\mathbb{H}^3(-1)$  of surfaces in hyperbolic 3-space  $\mathbb{H}^3(-1)$  in [4].

The anti-de Sitter (abbreviated: AdS) 3-space  $\mathbb{H}_1^3(-1)$  is a Lorentzian 3-manifold of sectional curvature  $-1$  that can be realized as the hyperquadric in  $\mathbb{E}_2^4$ :

$$\mathbb{H}_1^3(-1) := \{(x_0, x_1, x_2, x_3) \in \mathbb{E}_2^4 : -(x_0)^2 - (x_1)^2 + (x_2)^2 + (x_3)^2 = -1\}.$$

Let  $M$  be a connected orientable 2-manifold and  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  an immersion. The immersion  $\varphi$  is said to be *timelike* if the induced metric  $I$  on  $M$  is Lorentzian. The induced Lorentzian metric  $I$  determines a Lorentzian conformal structure  $\mathcal{C}_I$  on  $M$ .

Let  $(x, y)$  be a Lorentz isothermal coordinate system with respect to the conformal structure  $\mathcal{C}_I$ . Then the first fundamental form

$$I = \langle d\varphi, d\varphi \rangle$$

is given by the matrix  $I = e^\omega \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ . The first fundamental form is also written in terms of  $(x, y)$  as  $I = e^\omega \{-(dx)^2 + (dy)^2\}$ . Let  $u := x + y$  and  $v := -x + y$ . Then  $(u, v)$  defines a *null coordinate system* with respect to the conformal structure  $\mathcal{C}_I$ . The first fundamental form  $I$  is written in terms of  $(u, v)$  as

$$I = e^\omega du dv.$$

In terms of null coordinates  $u$  and  $v$ , the differential operators  $\frac{\partial}{\partial u}$  and  $\frac{\partial}{\partial v}$  are computed to be

$$\frac{\partial}{\partial u} = \frac{1}{2} \left( \frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial v} = \frac{1}{2} \left( -\frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right).$$

The conformality condition is equivalent to

$$\langle \varphi_u, \varphi_u \rangle = \langle \varphi_v, \varphi_v \rangle = 0, \quad \langle \varphi_u, \varphi_v \rangle = \frac{1}{2} e^\omega.$$

Let  $N$  be a unit normal vector field of  $M$ . Then

$$\langle N, N \rangle = 1, \quad \langle \varphi, N \rangle = \langle \varphi_u, N \rangle = \langle \varphi_v, N \rangle = 0.$$

The mean curvature  $H$  is given by  $H = 2e^{-\omega} \langle \varphi_{uv}, N \rangle$ . Let  $Q := \langle \varphi_{uu}, N \rangle$  and  $R := \langle \varphi_{vv}, N \rangle$ . Then the quadratic differentials  $Q^\sharp := Q du \otimes du$  and  $R^\sharp := R dv \otimes dv$  are called *Hopf pairs*<sup>2</sup> of  $M$ . The quadratic differential

$$\mathcal{Q} := Q du^2 + R dv^2 = Q^\sharp + R^\sharp$$

is called *Hopf differential*<sup>3</sup>. This differential is globally defined on the Lorentz surface  $(M, \mathcal{C}_I)$ . The second fundamental form  $II$  of  $M$  derived from  $N$  is defined by

$$II := -\langle d\varphi, dN \rangle$$

<sup>2</sup>In [6], [11], [13], the quadratic differentials  $Q^\sharp$  and  $R^\sharp$  are defined as Hopf differentials.

<sup>3</sup>The definition of Hopf differential  $\mathcal{Q}$  was suggested to the author by J. Inoguchi [12].

and it is given by the matrix

$$II = \begin{pmatrix} Q + R - He^\omega & Q - R \\ Q - R & Q + R + He^\omega \end{pmatrix}$$

with respect to Lorentz isothermal coordinate system  $(x, y)$ . The second fundamental form is related to Hopf differential  $\mathcal{Q}$  by

$$(1) \quad II = \mathcal{Q} + HI.$$

The *shape operator*  $\mathcal{S}$  of  $M$  derived from  $N$  is  $\mathcal{S} := -dN$ . The shape operator  $\mathcal{S}$  is related to  $II$  by

$$II(X, Y) = \langle \mathcal{S}X, Y \rangle$$

for all vector fields  $X, Y$  on  $M$ . The shape operator  $\mathcal{S}$  is also represented by the matrix  $II \cdot I^{-1}$ . The mean curvature  $H$  of  $M$  is

$$H = \frac{1}{2} \operatorname{tr} \mathcal{S} = \frac{1}{2} \operatorname{tr}(II \cdot I^{-1})$$

and the Gaußian curvature<sup>4</sup>  $K$  of  $M$  is

$$K := -1 + \det \mathcal{S} = -1 + \det(II \cdot I^{-1}).$$

The eigenvalues of  $\mathcal{S}$ , i.e., the solutions to the characteristic equation

$$\det(\mathcal{S} - \lambda \mathcal{I}) = 0, \quad \mathcal{I} = \text{identity of } TM$$

are called the principal curvatures. Since the metric  $I$  is indefinite, both principal curvatures may be nonreal complex numbers. The mean curvature  $H$  is the mean of the two principal curvatures and the Gaußian curvature  $K$  is the product of the two principal curvatures minus one.

A point  $p \in M$  is said to be an umbilic point if  $II$  is proportional to  $I$  at  $p$ . Equivalently,  $p$  is an umbilic point if and only if the two principal curvatures at  $p$  are the same real number and the corresponding eigenspace is 2-dimensional. A timelike surface is said to be a totally umbilic if all the points are umbilical. The formula (1) implies that  $p \in M$  is an umbilic point if and only if  $\mathcal{Q}(p) = 0$ , i.e.,  $p \in M$  is a common zero of Hopf pairs  $Q$  and  $R$ .

The Gauß equation which describes a relationship between  $K$ ,  $H$ ,  $Q$  and  $R$  takes the following form:

$$(2) \quad H^2 - K - 1 = 4e^{-2\omega}QR.$$

Note that the condition  $QR = 0$  does not imply the condition  $Q = R = 0$  (See [19]).

Let  $M$  be a simply-connected open and orientable 2-manifold and  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  a timelike conformal immersion with unit normal vector field  $N$ . Then we can define an orthonormal frame field  $\mathcal{F}$  along  $\varphi$  by

$$(3) \quad \mathcal{F} = (\varphi, e^{-\frac{\omega}{2}}\varphi_x, e^{-\frac{\omega}{2}}\varphi_y, N) : M \longrightarrow \mathcal{O}^{++}(2, 2),$$

---

<sup>4</sup>This can be easily computed from the Gauß equation (66)

where  $O^{++}(2, 2)$  denotes the identity component of the Lorentz group

$$O(2, 2) = \{\mathcal{A} \in GL_4\mathbb{R} : \langle \mathcal{A}\mathbf{u}, \mathcal{A}\mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle, \mathbf{u}, \mathbf{v} \in \mathbb{E}_2^4\}.$$

In terms of null coordinates  $(u, v)$ ,  $\mathcal{F}$  is defined by

$$(4) \quad \mathcal{F} = (\varphi, e^{-\frac{u}{2}}(\varphi_u - \varphi_v), e^{-\frac{v}{2}}(\varphi_u + \varphi_v), N) : M \longrightarrow O^{++}(2, 2).$$

The semi-Euclidean 4-space  $\mathbb{E}_2^4$  is identified with the linear space  $M_2\mathbb{R}$  of all  $2 \times 2$  real matrices via the correspondence

$$\mathbf{u} = (x_0, x_1, x_2, x_3) \longleftrightarrow \begin{pmatrix} x_0 + x_3 & x_1 + x_2 \\ -x_1 + x_2 & x_0 - x_3 \end{pmatrix}.$$

The scalar product of  $\mathbb{E}_2^4$  corresponds to the scalar product

$$(5) \quad \langle \mathbf{u}, \mathbf{v} \rangle = \frac{1}{2} \{ \text{tr}(\mathbf{u}\mathbf{v}) - \text{tr}(\mathbf{u})\text{tr}(\mathbf{v}) \}, \mathbf{u}, \mathbf{v} \in M_2\mathbb{R}.$$

Note that  $\langle \mathbf{u}, \mathbf{u} \rangle = -\det \mathbf{u}$ . The standard basis  $e_0, e_1, e_2, e_3$  for  $\mathbb{E}_2^4$  is identified with the matrices

$$\mathbf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \mathbf{i} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \mathbf{j}' = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \mathbf{k}' = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

i.e.,

$$(6) \quad (x_0, x_1, x_2, x_3) \longleftrightarrow x_0\mathbf{1} + x_1\mathbf{i} + x_2\mathbf{j}' + x_3\mathbf{k}' = \begin{pmatrix} x_0 + x_3 & x_1 + x_2 \\ -x_1 + x_2 & x_0 - x_3 \end{pmatrix}.$$

Note that the  $2 \times 2$  matrices  $x_0\mathbf{1} + x_1\mathbf{i} + x_2\mathbf{j}' + x_3\mathbf{k}'$  form the algebra  $\mathbb{H}'$  of *split-quaternions*. (For more details, see, for example, [13].) Under the identification (6), the group  $G$  of timelike unit vectors corresponds to a special linear group

$$SL_2\mathbb{R} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2\mathbb{R} : ad - bc = 1 \right\}.$$

The metric of  $G$  induced by the scalar product (5) is a bi-invariant Lorentz metric of constant curvature  $-1$ . Hence,  $G$  is identified with  $\mathbb{H}_1^3(-1)$ .

### 3. CARTAN'S FORMALISM

Let  $\{e_\alpha : \alpha = 0, 1, 2, 3\}$  be a frame field of  $\mathbb{E}_2^4$ , i.e.,  $\{e_\alpha(p) : \alpha = 0, 1, 2, 3\}$  is a basis for the tangent space  $T_p\mathbb{E}_2^4$  at each  $p \in \mathbb{E}_2^4$ . Denote by  $\langle \cdot, \cdot \rangle_p$  the scalar product on the tangent space  $T_p\mathbb{E}_2^4$ ,  $p \in \mathbb{E}_2^4$ . Then

$$\langle e_\alpha, e_\beta \rangle = \begin{cases} -1 & \text{if } \alpha = \beta = 0 \text{ or } 1, \\ 0 & \text{if } \alpha \neq \beta, \\ 1 & \text{if } \alpha = \beta = 2 \text{ or } 3. \end{cases}$$

There exist unique connection 1-forms  $\{\omega_\alpha^\beta : \alpha, \beta = 0, 1, 2, 3\}$  such that

$$(7) \quad de_\alpha = \omega_\alpha^\beta e_\beta.$$

We use the index range  $1 \leq i, j, k \leq 3$  and denote by  $\omega^i$  the connection form  $\omega_0^i$ . Then the equation (7) can be written

$$(8) \quad de_0 = \omega^i e_i,$$

$$(9) \quad de_1 = \omega_1^0 e_0 + \omega_1^2 e_2 + \omega_1^3 e_3,$$

$$(10) \quad de_i = \omega_i^0 e_0 + \omega_i^j e_j, \quad i = 2, 3.$$

The connection 1-forms  $\{\omega_\alpha^\beta : \alpha, \beta = 0, 1, 2, 3\}$  satisfy:

$$\begin{aligned} \omega_0^1 &= -\omega_1^0, \quad \omega_\alpha^\alpha = 0, \quad \alpha = 0, 1, 2, 3, \\ \omega_1^i &= \omega_i^1, \quad \omega_0^i = \omega_i^0, \quad i = 2, 3, \\ \omega_i^j &= -\omega_j^i, \quad i, j = 2, 3. \end{aligned}$$

Differentiating the equation (8) we get the *first structure equation*:

$$(11) \quad d\omega^i = -\omega_j^i \wedge \omega^j.$$

Differentiating this first structure equation (11) we get the *second structure equation*:

$$(12) \quad d\omega_j^i = -\omega_k^i \wedge \omega_j^k.$$

For the frame field  $\mathcal{F}$  of timelike immersion  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$ , we have

$$(13) \quad d\omega^1 = \omega^2 \wedge \omega_2^1 \quad (\text{The First Structure Equations})$$

$$(14) \quad d\omega^2 = \omega^1 \wedge \omega_2^1$$

$$(15) \quad 0 = \omega^1 \wedge \omega_3^1 - \omega^2 \wedge \omega_3^2 \quad (\text{Symmetry Equation})$$

$$(16) \quad d\omega_2^1 = -\omega^1 \wedge \omega^2 + \omega_3^1 \wedge \omega_3^2 \quad (\text{Gauß Equation})$$

$$(17) \quad d\omega_3^1 = -\omega_2^1 \wedge \omega_3^2 \quad (\text{Mainardi-Codazzi Equations})$$

$$(18) \quad d\omega_3^2 = -\omega_2^2 \wedge \omega_3^1.$$

**Proposition 1.** *Let  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  be a timelike immersion. If  $\{e_1 = e^{-\frac{\omega}{2}} \varphi_x, e_2 = e^{-\frac{\omega}{2}} \varphi_y, e_3 = N\}$  forms an adapted frame field along  $\varphi$ , then the Gaußian curvature  $K$  and mean curvature  $H$  of  $\varphi$  satisfy the following equations:*

$$(19) \quad \omega_3^1 \wedge \omega_3^2 = (K + 1) \omega^1 \wedge \omega^2,$$

$$(20) \quad \omega_3^1 \wedge \omega^2 + \omega^1 \wedge \omega_3^2 = -2H \omega^1 \wedge \omega^2.$$

*Proof.* From the symmetry equation (15), we see that there exist smooth functions  $h_{ij}$ ,  $i, j = 1, 2$  such that

$$\begin{pmatrix} \omega_3^1 \\ \omega_3^2 \end{pmatrix} = \begin{pmatrix} -h_{11} & h_{12} \\ h_{21} & -h_{22} \end{pmatrix} \begin{pmatrix} \omega^1 \\ \omega^2 \end{pmatrix} \quad \text{and} \quad h_{12} = -h_{21}.$$

Note that  $\omega^1$  and  $\omega^2$  are the dual 1-forms of  $e_1$  and  $e_2$ , resp., and so

$$\begin{aligned}\omega_3^1 \wedge \omega_3^2 &= (h_{11}h_{22} + h_{12}^2)\omega^1 \wedge \omega^2 \\ &= (K + 1)\omega^1 \wedge \omega^2,\end{aligned}$$

where  $K$  is the Gaußian curvature of  $\varphi$ . Thus, the Gaußian equation (16) can be written as  $d\omega_2^1 = K\omega^1 \wedge \omega^2$ . The mean curvature  $H$  of  $\varphi$  is  $\frac{h_{11} + h_{22}}{2}$ . Hence,

$$\begin{aligned}\omega_3^1 \wedge \omega^2 + \omega^1 \wedge \omega_3^2 &= -(h_{11} + h_{22})\omega^1 \wedge \omega^2 \\ &= -2H\omega^1 \wedge \omega^2.\end{aligned}$$

□

#### 4. LIE GROUP ACTIONS $\mu$ AND $\nu$ ON $\mathbb{E}_2^4$

The Lie group  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  acts isometrically on  $\mathbb{E}_2^4$  via the group action:

$$\mu : (\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}) \times \mathbb{E}_2^4 \longrightarrow \mathbb{E}_2^4; \mu(g_1, g_2)\mathbf{u} = g_1\mathbf{u}g_2^t.$$

This action is transitive on  $\mathbb{H}_1^3(-1)$ . The isotropy subgroup of  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  at  $\mathbf{1}$  is  $\mathcal{K} = \{(g, (g^{-1})^t) : g \in \mathrm{SL}_2\mathbb{R}\}$  and  $\mathbb{H}_1^3(-1)$  is represented as the Lorentzian symmetric space  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}/\mathcal{K}$ . The natural projection  $\pi_\mu : \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R} \longrightarrow \mathbb{H}_1^3(-1)$  is given explicitly by  $\pi_\mu(g_1, g_2) = g_1g_2^t$ .

The Lie group  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  also acts isometrically on  $\mathbb{E}_2^4$  via the diagonal action:

$$\nu : (\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}) \times \mathbb{E}_2^4 \longrightarrow \mathbb{E}_2^4; \nu(g_1, g_2)\mathbf{u} = g_1\mathbf{u}g_2^{-1}.$$

This action is also transitive on  $\mathbb{H}_1^3(-1)$ . The isotropy subgroup of  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  at  $\mathbf{1}$  is the diagonal subgroup  $\Delta$  of  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$ , that is,  $\Delta = \{(g, g) : g \in \mathrm{SL}_2\mathbb{R}\}$  and  $\mathbb{H}_1^3(-1)$  is also represented by  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}/\Delta$  as a Lorentzian symmetric space. The natural projection  $\pi_\nu : \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R} \longrightarrow \mathbb{H}_1^3(-1)$  is given explicitly by

$$\pi_\nu(g_1, g_2) = g_1g_2^{-1}, (g_1, g_2) \in \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}.$$

Moreover,  $\mathrm{SL}_2\mathbb{R}$  acts isometrically on  $\mathbb{E}_1^3$  via the Ad-action:

$$\mathrm{Ad} : \mathrm{SL}_2\mathbb{R} \times \mathbb{E}_1^3 \longrightarrow \mathbb{E}_1^3; \mathrm{Ad}(g)\mathbf{u} = g\mathbf{u}g^{-1}, g \in \mathrm{SL}_2\mathbb{R}, \mathbf{u} \in \mathbb{E}_1^3.$$

The actions  $\mu$  and  $\nu$  both induces a double covering  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R} \longrightarrow \mathrm{O}^{++}(2, 2)$  of the Lorentz group  $\mathrm{O}^{++}(2, 2)$ .

*Remark 1.* In [10], J. Q. Hong used the group action  $\mu$  to study a Bryant type representation formula for timelike cmc 1 surfaces in  $\mathbb{H}_1^3(-1)$ . In [1], R. Aiyama and K. Akutagawa also used the action  $\mu$  to study Kenmotsu-Bryant type representation formula for spacelike cmc surfaces in  $\mathbb{H}_1^3(-1)$ . In this paper, we use both actions  $\mu$  and  $\nu$ .

The frame field  $\{e_\alpha : \alpha = 0, 1, 2, 3\}$  can be parametrized by the Lie group  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  via the Lie group action  $\mu$ : for each  $g = (g_1, g_2) \in \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$ ,

$$\begin{aligned} e_0(g) &:= \mu(g)(\mathbf{1}) = g_1 \mathbf{1} g_2^t, \\ e_1(g) &:= \mu(g)(\mathbf{i}) = g_1 \mathbf{i} g_2^t, \\ e_2(g) &:= \mu(g)(\mathbf{j}') = g_1 \mathbf{j}' g_2^t, \\ e_3(g) &:= \mu(g)(\mathbf{k}') = g_1 \mathbf{k}' g_2^t. \end{aligned}$$

The frame field  $\{e_\alpha : \alpha = 0, 1, 2, 3\}$  can also be parametrized by the Lie group  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  via the Lie group action  $\nu$ : for each  $g = (g_1, g_2) \in \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$ ,

$$\begin{aligned} e_0(g) &:= \nu(g)(\mathbf{1}) = g_1 \mathbf{1} g_2^{-1}, \\ e_1(g) &:= \nu(g)(\mathbf{i}) = g_1 \mathbf{i} g_2^{-1}, \\ e_2(g) &:= \nu(g)(\mathbf{j}') = g_1 \mathbf{j}' g_2^{-1}, \\ e_3(g) &:= \nu(g)(\mathbf{k}') = g_1 \mathbf{k}' g_2^{-1}. \end{aligned}$$

We need the following two equations in order to do some differential geometric computations in Sections 6 and 7.

**Lemma 2.** *If the frame field  $\{e_\alpha : \alpha = 0, 1, 2, 3\}$  is parametrized by  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  via the action  $\mu$ , then the pull back  $g^{-1}dg$  of Maurer-Cartan form  $\Omega = (\omega_\alpha^\beta)$  can be written as the following equation in the Lie algebra  $\mathfrak{sl}_2\mathbb{R} \oplus \mathfrak{sl}_2\mathbb{R}$ :*

$$g^{-1}dg = g_1^{-1}dg_1 \oplus g_2^{-1}dg_2,$$

where

$$(21) \quad g_1^{-1}dg_1 = \frac{1}{2} \begin{pmatrix} \omega^3 + \omega_2^1 & \omega^1 + \omega^2 - \omega_3^1 - \omega_3^2 \\ -\omega^1 + \omega^2 - \omega_3^1 + \omega_3^2 & -\omega^3 - \omega_2^1 \end{pmatrix}$$

and

$$(22) \quad g_2^{-1}dg_2 = \frac{1}{2} \begin{pmatrix} \omega^3 - \omega_2^1 & -\omega^1 + \omega^2 + \omega_3^1 - \omega_3^2 \\ \omega^1 + \omega^2 + \omega_3^1 + \omega_3^2 & -\omega^3 + \omega_2^1 \end{pmatrix}.$$

*Proof.* For simplicity, let  $\sigma_0 := \mathbf{1}, \sigma_1 := \mathbf{i}, \sigma_2 := \mathbf{j}', \sigma_3 := \mathbf{k}'$ . By applying the chain rule,

$$\begin{aligned} de_\alpha(g) &= d(g_1 \sigma_\alpha g_2^t) \\ &= (dg_1) \sigma_\alpha g_2^t + g_1 \sigma_\alpha dg_2^t \\ &= g_1 \{g_1^{-1}(dg_1) \sigma_\alpha + \sigma_\alpha (g_2^{-1}dg_2)^t\} g_2^t. \end{aligned}$$

On the other hand,

$$de_\alpha = \omega_\alpha^\beta e_\beta = \omega_\alpha^\beta g_1 \sigma_\beta g_2.$$

Hence, we have the equation

$$(g_1^{-1}dg_1) \sigma_\alpha + \sigma_\alpha (g_2^{-1}dg_2)^t = \omega_\alpha^\beta \sigma_\beta.$$

The equations (21) and (22) follow from this equation.  $\square$

**Lemma 3.** *If the frame field  $\{e_\alpha : \alpha = 0, 1, 2, 3\}$  is parametrized by  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  via the action  $\nu$ , then the pull back  $g^{-1}dg$  of Maurer-Cartan form  $\Omega = (\omega_\alpha^\beta)$  can be written as the following equation in the Lie algebra  $\mathfrak{sl}_2\mathbb{R} \oplus \mathfrak{sl}_2\mathbb{R}$ :*

$$g^{-1}dg = g_1^{-1}dg_1 \oplus (dg_2^{-1})g_2,$$

where

$$(23) \quad g_1^{-1}dg_1 = \frac{1}{2} \begin{pmatrix} \omega^3 + \omega_2^1 & \omega^1 + \omega^2 - \omega_3^1 - \omega_3^2 \\ -\omega^1 + \omega^2 - \omega_3^1 + \omega_3^2 & -\omega^3 - \omega_2^1 \end{pmatrix}$$

and

$$(24) \quad (dg_2^{-1})g_2 = \frac{1}{2} \begin{pmatrix} \omega^3 - \omega_2^1 & \omega^1 + \omega^2 + \omega_3^1 + \omega_3^2 \\ -\omega^1 + \omega^2 + \omega_3^1 - \omega_3^2 & -\omega^3 + \omega_2^1 \end{pmatrix}.$$

*Proof.* Similar to the proof of Lemma 2, we get the equation

$$(g_1^{-1}dg_1)\sigma_\alpha + \sigma_\alpha(dg_2^{-1})g_2 = \omega_\alpha^\beta\sigma_\beta$$

and the equations (23) and (24) then follow.  $\square$

## 5. TIMELIKE cmc SURFACES IN ADS 3-SPACE $\mathbb{H}_1^3(-1)$ AND INTEGRABLE SYSTEMS

Let  $M$  be a simply-connected open and orientable 2-manifold and  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  a timelike conformal immersion.

By using a double covering induced by the group action  $\mu$ , we can find lift  $\Phi = (\Phi_1, \Phi_2)$  (called a coordinate frame) of  $\mathcal{F}$  to  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$ :

$$\mu(\Phi)(\mathbf{1}, \mathbf{i}, \mathbf{j}', \mathbf{k}') = \mathcal{F}.$$

That is, the lifted framing  $\Phi = (\Phi_1, \Phi_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  satisfies<sup>5</sup>

$$\begin{aligned} \mu(\Phi)(\mathbf{1}) &= \Phi_1 \mathbf{1} \Phi_2^t = \varphi, \\ \mu(\Phi)(\mathbf{i}) &= \Phi_1 \mathbf{i} \Phi_2^t = e^{-\frac{\omega}{2}} \varphi_x, \\ \mu(\Phi)(\mathbf{j}') &= \Phi_1 \mathbf{j}' \Phi_2^t = e^{-\frac{\omega}{2}} \varphi_y, \\ \mu(\Phi)(\mathbf{k}') &= \Phi_1 \mathbf{k}' \Phi_2^t = N. \end{aligned}$$

Then

$$(25) \quad \varphi_u = e^{\frac{\omega}{2}} \Phi_1 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \Phi_2^t$$

and

$$(26) \quad \varphi_v = e^{\frac{\omega}{2}} \Phi_1 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \Phi_2^t.$$

---

<sup>5</sup>Here, we use the same  $\mu$  for both Lie group action and group representation.

Similarly, by using a double covering induced by the group action  $\nu$ , we can find lift  $\Psi = (\Psi_1, \Psi_2)$  of  $\mathcal{F}$  to  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$ :

$$\nu(F)(\mathbf{1}, \mathbf{i}, \mathbf{j}', \mathbf{k}') = \mathcal{F}.$$

The lifted framing  $\Psi = (\Psi_1, \Psi_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  satisfies

$$\begin{aligned}\nu(\Psi)(\mathbf{1}) &= \Psi_1 \mathbf{1} \Psi_2^{-1} = \varphi, \\ \nu(\Psi)(\mathbf{i}) &= \Psi_1 \mathbf{i} \Psi_2^{-1} = e^{-\frac{\omega}{2}} \varphi_x, \\ \nu(\Psi)(\mathbf{j}') &= \Psi_1 \mathbf{j}' \Psi_2^{-1} = e^{-\frac{\omega}{2}} \varphi_y, \\ \nu(\Psi)(\mathbf{k}') &= \Psi_1 \mathbf{k}' \Psi_3^{-1} = N.\end{aligned}$$

Then

$$(27) \quad \varphi_u = e^{\frac{\omega}{2}} \Psi_1 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \Psi_2^{-1}$$

and

$$(28) \quad \varphi_v = e^{\frac{\omega}{2}} \Psi_1 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \Psi_2^{-1}.$$

Let  $\mathfrak{s} := (\varphi, \varphi_u, \varphi_v, N)$ . Then  $\mathfrak{s}$  defines a moving frame on the immersed surface  $\varphi$  and satisfy the following Gauß-Weingarten equations:

$$(29) \quad \mathfrak{s}_u = \mathfrak{s}\mathcal{U}, \quad \mathfrak{s}_v = \mathfrak{s}\mathcal{V},$$

where

$$(30) \quad \mathcal{U} = \begin{pmatrix} 0 & 0 & \frac{1}{2}e^\omega & 0 \\ 1 & \omega_u & 0 & -H \\ 0 & 0 & 0 & -2Qe^{-\omega} \\ 0 & Q & \frac{1}{2}e^\omega H & 0 \end{pmatrix}, \quad \mathcal{V} = \begin{pmatrix} 0 & \frac{1}{2}e^\omega & 0 & 0 \\ 0 & 0 & 0 & -2Re^{-\omega} \\ 1 & 0 & \omega_v & -H \\ 0 & \frac{1}{2}e^\omega H & R & 0 \end{pmatrix}.$$

The integrability condition of the Gauß-Weingarten equation is the Gauß-Mainardi-Codazzi equation

$$\mathcal{V}_u - \mathcal{U}_v + [\mathcal{U}, \mathcal{V}] = 0.$$

This Gauß-Mainardi-Codazzi equation is equivalent to

$$(31) \quad \omega_{uv} + \frac{1}{2}e^\omega(H^2 - 1) - 2QRe^{-\omega} = 0,$$

$$(32) \quad H_u = 2e^{-\omega}Q_v, \quad H_v = 2e^{-\omega}R_u.$$

*Remark 2.* From the equations (32), we see that a timelike surface  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  has constant mean curvature if and only if  $R_u = Q_v = 0$ . In this case,  $R$  is said to be *Lorentz antiholomorphic* and  $Q$  is said to be *Lorentz holomorphic*, respectively.

Each component framing  $\Phi_1$  and  $\Phi_2$  of  $\Phi$  satisfy the following Lax equations:

$$\begin{aligned} (\Phi_1)_u &= \Phi_1 \mathcal{U}_1, & (\Phi_1)_v &= \Phi_1 \mathcal{V}_1; \\ (\Phi_2)_u &= \Phi_2 \mathcal{U}_2, & (\Phi_2)_v &= \Phi_2 \mathcal{V}_2, \end{aligned}$$

where

$$(33) \quad \mathcal{U}_1 = \begin{pmatrix} \frac{\omega_u}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H+1) \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix},$$

$$(34) \quad \mathcal{V}_1 = \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}}R \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H-1) & \frac{\omega_v}{4} \end{pmatrix},$$

$$(35) \quad \mathcal{U}_2 = \begin{pmatrix} -\frac{\omega_u}{4} & e^{-\frac{\omega}{2}}Q \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H-1) & \frac{\omega_u}{4} \end{pmatrix},$$

$$(36) \quad \mathcal{V}_2 = \begin{pmatrix} \frac{\omega_v}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H+1) \\ -e^{-\frac{\omega}{2}}R & -\frac{\omega_v}{4} \end{pmatrix}.$$

The compatibility conditions  $(\Phi_1)_{uv} = (\Phi_1)_{vu}$  and  $(\Phi_2)_{uv} = (\Phi_2)_{vu}$  give the Maurer-Cartan equations

$$(37) \quad (\mathcal{U}_1)_v - (\mathcal{V}_1)_u - [\mathcal{U}_1, \mathcal{V}_1] = 0$$

and

$$(38) \quad (\mathcal{U}_2)_v - (\mathcal{V}_2)_u - [\mathcal{U}_2, \mathcal{V}_2] = 0.$$

Each of these two Maurer-Cartan equations is also equivalent to the Gauß-Mainardi-Codazzi equations (31) and (32).

Each component framing  $\Psi_1$  and  $\Psi_2$  of  $\Psi$  satisfy the following Lax equations:

$$\begin{aligned} (\Psi_1)_u &= \Psi_1 \mathcal{U}_1, & (\Psi_1)_v &= \Psi_1 \mathcal{V}_1; \\ (\Psi_2)_u &= \Psi_2 \mathcal{U}_2, & (\Psi_2)_v &= \Psi_2 \mathcal{V}_2, \end{aligned}$$

where

$$(39) \quad \mathcal{U}_1 = \begin{pmatrix} \frac{\omega_u}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H+1) \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix},$$

$$(40) \quad \mathcal{V}_1 = \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}}R \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H-1) & \frac{\omega_v}{4} \end{pmatrix},$$

$$(41) \quad \mathcal{U}_2 = \begin{pmatrix} \frac{\omega_u}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H-1) \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix},$$

$$(42) \quad \mathcal{V}_2 = \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}}R \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H+1) & \frac{\omega_v}{4} \end{pmatrix}.$$

The compatibility conditions  $(\Psi_1)_{uv} = (\Psi_1)_{vu}$  and  $(\Psi_2)_{uv} = (\Psi_2)_{vu}$  give the Maurer-Cartan equations

$$(43) \quad (\mathcal{U}_1)_v - (\mathcal{V}_1)_u - [\mathcal{U}_1, \mathcal{V}_1] = 0$$

and

$$(44) \quad (\mathcal{U}_2)_v - (\mathcal{V}_2)_u - [\mathcal{U}_2, \mathcal{V}_2] = 0.$$

Again, each of these two Maurer-Cartan equations is equivalent to the Gauß-Mainardi-Codazzi equations (31) and (32).

We now have the following representation formulae for timelike cmc surfaces in  $\mathbb{H}_1^3(-1)$ .

**Theorem 4.** *Let  $M$  be a simply-connected region in Minkowski plane  $\mathbb{E}_1^2 = (\mathbb{R}^2(u, v), dudv)$ .*

- (1) *Let  $\Phi = (\Phi_1, \Phi_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  be a solution to the following Lax equations:*

$$(45) \quad \begin{aligned} (\Phi_1)_u &= \Phi_1 \mathcal{U}_1, & (\Phi_1)_v &= \Phi_1 \mathcal{V}_1; \\ (\Phi_2)_u &= \Phi_2 \mathcal{U}_2, & (\Phi_2)_v &= \Phi_2 \mathcal{V}_2, \end{aligned}$$

where

$$\begin{aligned} \mathcal{U}_1 &= \begin{pmatrix} \frac{\omega_u}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H+1) \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_1 &= \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}}R \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H-1) & \frac{\omega_v}{4} \end{pmatrix}, \\ \mathcal{U}_2 &= \begin{pmatrix} -\frac{\omega_u}{4} & e^{-\frac{\omega}{2}}Q \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H-1) & \frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_2 &= \begin{pmatrix} \frac{\omega_v}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H+1) \\ -e^{-\frac{\omega}{2}}R & -\frac{\omega_v}{4} \end{pmatrix}. \end{aligned}$$

Then  $\varphi := \mu(\Phi)(\mathbf{1}) = \Phi_1 \Phi_2^t : M \longrightarrow \mathbb{H}_1^3(-1)$  defines a timelike cmc  $H$  immersion into  $\mathbb{H}_1^3(-1)$ .

- (2) *Let  $\Psi = (\Psi_1, \Psi_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  be a solution to the following Lax equations:*

$$(46) \quad \begin{aligned} (\Psi_1)_u &= \Psi_1 \mathcal{U}_1, & (\Psi_1)_v &= \Psi_1 \mathcal{V}_1; \\ (\Psi_2)_u &= \Psi_2 \mathcal{U}_2, & (\Psi_2)_v &= \Psi_2 \mathcal{V}_2, \end{aligned}$$

where

$$\begin{aligned} \mathcal{U}_1 &= \begin{pmatrix} \frac{\omega_u}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H+1) \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_1 &= \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}}R \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H-1) & \frac{\omega_v}{4} \end{pmatrix}, \\ \mathcal{U}_2 &= \begin{pmatrix} \frac{\omega_u}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H-1) \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_2 &= \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}}R \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H+1) & \frac{\omega_v}{4} \end{pmatrix}. \end{aligned}$$

Then  $\psi := \nu(\Psi)(\mathbf{1}) = \Psi_1 \Psi_2^{-1} : M \longrightarrow \mathbb{H}_1^3(-1)$  defines a timelike cmc  $H$  immersion into  $\mathbb{H}_1^3(-1)$ .

Let  $H_e$ ,  $H_s$ ,  $H_h$  be the constant mean curvatures of timelike surfaces in Minkowski 3-space  $\mathbb{E}_1^3$ , de Sitter 3-space  $\mathbb{S}_1^3(1)$  and anti-de Sitter 3-space  $\mathbb{H}_1^3(-1)$ , resp. Then these timelike cmc surfaces in each space-form satisfy

the following Gauß-Mainardi-Codazzi equations:

$$\begin{aligned} \omega_{uv} + \frac{1}{2}H_e^2 e^\omega - 2QR e^{-\omega} &= 0 && \text{Lorentz 3-space } \mathbb{E}_1^3 \text{ case,} \\ \omega_{uv} + \frac{1}{2}(H_s^2 + 1)e^\omega - 2QR e^{-\omega} &= 0 && \text{de Sitter 3-space } \mathbb{S}_1^3(1) \text{ case,} \\ \omega_{uv} + \frac{1}{2}(H_h^2 - 1)e^\omega - 2QR e^{-\omega} &= 0 && \text{anti-de Sitter 3-space } \mathbb{H}_1^3(-1) \text{ case.} \end{aligned}$$

By comparing these Gauß-Mainardi-Codazzi equations, we can deduce the *Lawson-Guichard correspondence* between timelike cmc  $H_e$  surfaces in  $\mathbb{E}_1^3$ , timelike cmc  $H_s$  surfaces in  $\mathbb{S}_1^3(1)$  and timelike cmc  $H_h$  surfaces in  $\mathbb{H}_1^3(-1)$ , which satisfy the same Gauß-Mainardi-Codazzi equation. Such cmc surfaces are called *cousins* of each other. In particular, we see that there is a bijective correspondence between timelike minimal surfaces ( $H = 0$ ) in  $\mathbb{E}_1^3$  and timelike cmc  $\pm 1$  in  $\mathbb{H}_1^3(-1)$ . For this reason, we are mainly interested in timelike cmc  $\pm 1$  surfaces in  $\mathbb{H}_1^3(-1)$ . If  $H = \pm 1$  then the Gauß-Mainardi-Codazzi equations (31) become

$$(47) \quad \begin{cases} \omega_{uv} - 2e^{-\omega}RQ = 0, \\ R_u = Q_v = 0. \end{cases}$$

For more details about the Lawson-Guichard correspondence, please see the appendix I (section 14).

*Remark 3.* Note that one can normalize the Hopf pairs  $Q$  and  $R$  if  $M$  has real distinct principal curvatures or imaginary principal curvatures everywhere. For example,  $Q = \pm R = 1$  reduces the Gauß-Mainardi-Codazzi equations (31) to the Liouville equation  $\omega_{uv} = \mp 2e^{-\omega}$ .

Since the sign of  $H$  depends upon the orientation of a surface (i.e., the orientation of the unit normal vector field  $N$ ), hereafter we consider only  $H = 1$  case.

**Corollary 5.** *Let  $M$  be a simply-connected 2-manifold.*

- (1) *Let  $\Phi = (\Phi_1, \Phi_2) : M \longrightarrow \text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  be solutions to the following Lax equations:*

$$\begin{aligned} (\Phi_1)_u &= \Phi_1 \mathcal{U}_1, & (\Phi_1)_v &= \Phi_1 \mathcal{V}_1; \\ (\Phi_2)_u &= \Phi_2 \mathcal{U}_2, & (\Phi_2)_v &= \Phi_2 \mathcal{V}_2, \end{aligned}$$

where

$$\begin{aligned} \mathcal{U}_1 &= \begin{pmatrix} \frac{\omega_u}{4} & e^{\frac{\omega}{2}} \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_1 &= \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}}R \\ 0 & \frac{\omega_v}{4} \end{pmatrix}, \\ \mathcal{U}_2 &= \begin{pmatrix} -\frac{\omega_u}{4} & e^{-\frac{\omega}{2}}Q \\ 0 & \frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_2 &= \begin{pmatrix} \frac{\omega_v}{4} & e^{\frac{\omega}{2}} \\ -e^{-\frac{\omega}{2}}R & -\frac{\omega_v}{4} \end{pmatrix}. \end{aligned}$$

Then  $\varphi := \Phi_1 \Phi_2^t : M \longrightarrow \mathbb{H}_1^3(-1)$  defines a timelike cmc 1 immersion into  $\mathbb{H}_1^3(-1)$ .

(2) Let  $\Psi = (\Psi_1, \Psi_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  be solutions to the following Lax equations:

$$\begin{aligned} (\Psi_1)_u &= \Psi_1 \mathcal{U}_1, & (\Psi_1)_v &= \Psi_1 \mathcal{V}_1; \\ (\Psi_2)_u &= \Psi_2 \mathcal{U}_2, & (\Psi_2)_v &= \Psi_2 \mathcal{V}_2, \end{aligned}$$

where

$$\begin{aligned} \mathcal{U}_1 &= \begin{pmatrix} \frac{\omega_u}{4} & e^{\frac{\omega}{2}} \\ -e^{-\frac{\omega}{2}} Q & -\frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_1 &= \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}} R \\ 0 & \frac{\omega_v}{4} \end{pmatrix}, \\ \mathcal{U}_2 &= \begin{pmatrix} \frac{\omega_u}{4} & 0 \\ -e^{-\frac{\omega}{2}} Q & -\frac{\omega_u}{4} \end{pmatrix}, & \mathcal{V}_2 &= \begin{pmatrix} -\frac{\omega_v}{4} & e^{-\frac{\omega}{2}} R \\ e^{-\frac{\omega}{2}} & \frac{\omega_v}{4} \end{pmatrix}. \end{aligned}$$

Then  $\psi := \Psi_1 \Psi_2^{-1} : M \longrightarrow \mathbb{H}_1^3(-1)$  defines a timelike cmc 1 immersion into  $\mathbb{H}_1^3(-1)$ .

## 6. TIMELIKE SURFACES OF CONSTANT MEAN CURVATURE $\pm 1$ IN ADS 3-SPACE $\mathbb{H}_1^3(-1)$ VIA THE GROUP ACTION $\mu$

In [10], J. Q. Hong gave a Bryant type representation formula for timelike surfaces of constant mean curvature 1 in  $\mathbb{H}_1^3(-1)$ . In this section, we reproduce J. Q. Hong's Bryant type representation formula in a more general context.

Let  $F : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  be a lift of  $\mathcal{F}$  to  $\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  via the Lie group action  $\mu$ , i.e.,  $\mu(F)(\mathbf{1}, \mathbf{i}, \mathbf{j}', \mathbf{k}') = \mathcal{F}$ . Let  $\Omega := \Omega_1 \oplus \Omega_2 \in \mathfrak{sl}_2\mathbb{R} \oplus \mathfrak{sl}_2\mathbb{R}$ , where  $\Omega_i = F_i^{-1} dF_i \in \mathfrak{sl}_2\mathbb{R}$ ,  $i = 1, 2$ . The Gauß and Mainardi-Codazzi equations are equivalent to Maurer-Cartan equation

$$(48) \quad d\Omega + \Omega \wedge \Omega = 0,$$

which is the null curvature (integrability) condition of the Maurer-Cartan form  $\Omega$ . Let  $F^*$  denote the pull-back map  $F^* : (\mathfrak{sl}_2\mathbb{R})^* \oplus (\mathfrak{sl}_2\mathbb{R})^* \longrightarrow T^*M$ . The Maurer-Cartan forms  $\Omega_i = F_i^{-1} dF_i \in \mathfrak{sl}_2\mathbb{R}$ ,  $i = 1, 2$  can be written as the following equations:

$$(49) \quad F_1^{-1} dF_1 = \frac{1}{2} F^* \begin{pmatrix} \omega^3 + \omega_2^1 & \omega^1 + \omega^2 - \omega_3^1 - \omega_3^2 \\ -\omega_1 + \omega_2 - \omega_3^1 + \omega_3^2 & -\omega^3 - \omega_2^1 \end{pmatrix},$$

$$(50) \quad F_2^{-1} dF_2 = \frac{1}{2} F^* \begin{pmatrix} \omega^3 - \omega_2^1 & -\omega^1 + \omega^2 + \omega_3^1 - \omega_3^2 \\ \omega^1 + \omega^2 + \omega_3^1 + \omega_3^2 & -\omega^3 + \omega_2^1 \end{pmatrix}.$$

**Definition 1.** Let  $M$  be a 2-manifold. A map  $F : M \longrightarrow \mathrm{SL}_2\mathbb{R}$  is said to be null if  $F^*(\phi) = 0$ , or equivalently  $\det(F^{-1} dF) = 0$ , where  $\phi$  is the quadratic Cartan-Killing form  $\phi = -8 \det(g^{-1} dg)$ .

**Theorem 6** (A Bryant type representation formula for timelike cmc  $\pm 1$  Surfaces in  $\mathbb{H}_1^3(-1)$ ). Let  $M$  be an open 2-manifold and  $F = (F_1, F_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  an immersion such that

- (1)  $F_1$  is Lorentz holomorphic, i.e.,  $(F_1)_v = 0$  and  $F_2$  is Lorentz antiholomorphic, i.e.,  $(F_2)_u = 0$ ,

(2)  $F_1$  and  $F_2$  are null, i.e.,  $\det F_1^{-1}dF_1 = \det F_2^{-1}dF_2 = 0$ .

Then

$$(51) \quad \varphi := \mu(F)(\mathbf{1}) = F_1 F_2^t$$

is a smooth conformal timelike immersion into  $\mathbb{H}_1^3(-1)$  with cmc  $\pm 1$ . Conversely, let  $M$  be an oriented and simply-connected Lorentzian 2-manifold with globally defined null coordinates<sup>6</sup>. If  $\varphi : M \rightarrow \mathbb{H}_1^3(-1)$  is a smooth conformal timelike immersion with cmc  $\pm 1$ , then there exists an immersion  $F = (F_1, F_2) : M \rightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  such that  $F_1, F_2$  satisfy the conditions (1), (2), and  $\varphi = F_1 F_2^t$ .

*Proof.* Let  $\omega^+ = \omega^1 + \omega^2$ ,  $\omega^- = -\omega^1 + \omega^2$ ,  $\pi^+ = \omega_3^1 + \omega_3^2$ ,  $\pi^- = -\omega_3^1 + \omega_3^2$ . Also let

$$\begin{aligned} F^*(\omega^3 + \omega_2^1) &= 2\alpha_1, \quad F^*(\omega^+ - \pi^+) = 2\beta_1, \quad F^*(\omega^- + \pi^-) = 2\gamma_1 \\ F^*(\omega^3 - \omega_2^1) &= 2\alpha_2, \quad F^*(\omega^- - \pi^-) = 2\beta_2, \quad F^*(\omega^+ + \pi^+) = 2\gamma_2. \end{aligned}$$

Then the Maurer-Cartan equations (21) and (22) become

$$(52) \quad F_1^{-1}dF_1 = \frac{1}{2}F^* \begin{pmatrix} \omega^3 + \omega_2^1 & \omega^+ - \pi^+ \\ \omega^- + \pi^- & -\omega^3 - \omega_2^1 \end{pmatrix} = \begin{pmatrix} \alpha_1 & \beta_1 \\ \gamma_1 & -\alpha_1 \end{pmatrix},$$

$$(53) \quad F_2^{-1}dF_2 = \frac{1}{2}F^* \begin{pmatrix} \omega^3 - \omega_2^1 & \omega^- - \pi^- \\ \omega^+ + \pi^+ & -\omega^3 + \omega_2^1 \end{pmatrix} = \begin{pmatrix} \alpha_2 & \beta_2 \\ \gamma_2 & -\alpha_2 \end{pmatrix}.$$

Note that  $F^*(\omega^3) = \alpha_1 + \alpha_2$ ,  $F^*(\omega^+) = \beta_1 + \gamma_2$ ,  $F^*(\omega^-) = -(\beta_2 + \gamma_1)$ . Since  $\det F_1^{-1}dF_1 = \det F_2^{-1}dF_2 = 0$ ,  $\alpha_1^2 + \beta_1\gamma_1 = 0$  and  $\alpha_2^2 + \beta_2\gamma_2 = 0$ .

Denote by  $ds^2$  the metric in  $\mathbb{H}_1^3(-1)$  induced by the canonical semi-Riemannian metric in  $\mathbb{E}_2^4$ . If we regard  $e_0$  as a map  $e_0 : \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R} \rightarrow \mathbb{H}_1^3(-1)$  given by  $e_0(g) = \mu(g)(\mathbf{1})$ ,  $g \in \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$ , then

$$\begin{aligned} e_0^*(ds^2) &= \langle de_0, de_0 \rangle \\ &= \langle \omega^1 e_1 + \omega^2 e_2 + \omega^3 e_3, \omega^1 e_1 + \omega^2 e_2 + \omega^3 e_3 \rangle \\ &= -(\omega^1)^2 + (\omega^2)^2 + (\omega^3)^2 \end{aligned}$$

---

<sup>6</sup>In Lorentzian case, the so-called *Riemann Mapping Theorem* or *Köbe Uniformization Theorem* does not hold. So, globally defined null coordinates do not exist, in general, on a simply-connected Lorentzian 2-manifold.

defines an indefinite metric in the oriented orthonormal frame bundle of  $\mathbb{H}_1^3(-1)$ . Since  $\varphi = e_0 \circ F$ ,

$$\begin{aligned} ds_\varphi^2 &= \varphi^*(ds^2) \\ &= F^* \circ e_0^*(ds^2) \\ &= F^*((\omega^3)^2 + \{-(\omega^1)^2 + (\omega^2)^2\}) \\ &= F^*(\omega^3)^2 + F^*(\omega^+)F^*(\omega^-) \\ &= 2\alpha_1\alpha_2 + \beta_1\beta_2 + \gamma_1\gamma_2. \end{aligned}$$

Since  $F$  is an immersion, the last expression defines a metric.

We now show that for the immersion  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$ ,  $H \equiv 1$ . Let  $U \subset M$  be a simply-connected open set in which there exists a null coordinate system  $(u, v)$  such that  $ds_\varphi^2 = e^\omega du dv$  for some real-valued function  $\omega : U \longrightarrow \mathbb{R}$  defined on  $U$ . Clearly,  $M$  is covered by such open sets. Let  $\eta = e^{\frac{\omega}{2}} du$  and  $\xi = e^{\frac{\omega}{2}} dv$ . There exist functions  $A_1, A_2, B_1, B_2, C_1, C_2$  such that

$$\begin{aligned} F^*(\omega^3 + \omega_2^1) &= 2A_1\eta, \quad F^*(\omega^3 - \omega_2^1) = 2A_2\xi, \\ F^*(\omega^+ - \pi^+) &= 2B_1\eta, \quad F^*(\omega^- - \pi^-) = 2B_2\xi, \\ F^*(\omega^- + \pi^-) &= 2C_1\eta, \quad F^*(\omega^+ + \pi^+) = 2C_2\xi. \end{aligned}$$

We, then, have equations

$$A_1^2 + B_1C_1 = 0, \quad A_2 + B_2C_2 = 0, \quad \text{and} \quad 2A_1A_2 + B_1B_2 + C_1C_2 = 1.$$

In the open set  $U$ ,

$$\begin{aligned} ds_\varphi^2 &= (2A_1A_2 + B_1B_2 + C_1C_2)\eta\xi \\ &= \eta\xi. \end{aligned}$$

Since  $A_1^2 + B_1C_1 = 0$  and  $A_2 + B_2C_2 = 0$ , there exist smooth functions  $p_1, p_2, q_1, q_2$  defined in  $U$  (unique up to replacement by  $(-p_1, -q_1)$  and  $(-p_2, -q_2)$  respectively) such that

$$\begin{aligned} A_1 &= p_1q_1, & A_2 &= p_2q_2 \\ B_1 &= q_1^2, & B_2 &= q_2^2 \\ C_1 &= -p_1^2, & C_2 &= -p_2^2 \end{aligned}$$

and

$$2p_1q_1p_2q_2 + q_1^2q_2^2 + p_1^2p_2^2 = (p_1p_2 + q_1q_2)^2 = 1.$$

By the continuity of  $p_1p_2 + q_1q_2$ , either  $p_1p_2 + q_1q_2 = 1$  in  $U$  or  $p_1p_2 + q_1q_2 = -1$  in  $U$ . Without loss of generality, we may assume that  $p_1p_2 + q_1q_2 = 1$ . Now define a map  $h : U \longrightarrow \text{SL}_2\mathbb{R}$  by  $h = \begin{pmatrix} p_1 & -q_2 \\ q_1 & p_2 \end{pmatrix}$ . Then  $e_0(F_1h, F_2(h^{-1})^t) =$

$$e_0(F_1, F_2) = F_1 F_2^t.$$

$$\begin{aligned} (F_1 h)^{-1} d(F_1 h) &= (h^{-1} F_1^{-1})((dF_1)h + F_1 dh) \\ &= h^{-1}(F_1^{-1} dF_1)h + h^{-1} dh \\ &= h^{-1} \begin{pmatrix} p_1 q_1 & -p_1^2 \\ q_1^2 & -p_1 q_1 \end{pmatrix} \eta h + h^{-1} dh \\ &= \begin{pmatrix} p_2 dp_1 + q_2 dq_1 & -p_2 dq_2 + q_2 dp_2 - \eta \\ -q_1 dp_1 + p_1 dq_1 & q_1 dq_2 + p_1 dp_2 \end{pmatrix}. \end{aligned}$$

Similarly,

$$(F_2(h^{-1})^t)^{-1} d(F_2(h^{-1})^t) = \begin{pmatrix} p_1 dp_2 + q_1 dq_2 & -p_1 dq_1 + q_1 dp_1 - \xi \\ -q_2 dp_2 + p_2 dq_2 & q_2 dq_1 + p_2 dp_1 \end{pmatrix}.$$

It then follows that  $(Fh)^*(\omega^3) = 0$ ,  $(Fh)^*(\omega^+) = -\eta$  and  $(Fh)^*(\omega^-) = -\xi$ , i.e.,  $(F_1 h, F_2(h^{-1})^t) : U \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  is an oriented framing in  $U$  along the immersion  $\varphi = e_0 \circ F = e_0(F_1 h, F_2(h^{-1})^t) = F_1 F_2^t$ .

The 1-form  $\frac{1}{2}(Fh)^*(\omega^- + \pi^-) = p_1 dq_1 - q_1 dp_1$  can be written

$$p_1 dq_1 - q_1 dp_1 = \begin{cases} p_1^2 d\left(\frac{q_1}{p_1}\right) & \text{where } p_1 \neq 0 \\ -q_1^2 d\left(\frac{p_1}{q_1}\right) & \text{where } q_1 \neq 0 \end{cases}.$$

Hence,  $\frac{1}{2}(Fh)^*(\omega^- + \pi^-)$  is a 1-form of type  $(1, 0)$ , i.e., a multiple of the 1-form  $\eta$ . Similarly,  $(Fh)^*(\omega^+ + \pi^+)$  is a 1-form of type  $(0, 1)$ , i.e., a multiple of the 1-form  $\xi$ .

Since  $(Fh)^*(\omega^+) = -\eta$  and  $(Fh)^*(\omega^-) = -\xi$ , by the equation (20), one can easily see that:

- (1) If  $\varphi$  satisfies  $H = 1$  in  $U$ , then  $(Fh)^*(\omega^- + \pi^-)$  is a 1-form of type  $(1, 0)$  and  $(Fh)^*(\omega^+ + \pi^+)$  is a 1-form of type  $(0, 1)$ .
- (2) If  $(Fh)^*(\omega^- + \pi^-)$  is a 1-form of type  $(1, 0)$  or  $(Fh)^*(\omega^+ + \pi^+)$  is a 1-form of type  $(0, 1)$ , then  $\varphi$  satisfies  $H = 1$  in  $U$ .

Therefore, we conclude that  $H = 1$  in  $U$ .

Conversely, let  $M$  be an oriented and simply-connected Lorentzian 2-manifold with globally defined null coordinates  $(u, v)$ . Let  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  be a smooth conformal timelike immersion into  $\mathbb{H}_1^3(-1)$  with cmc 1. Then  $ds_\varphi^2 = e^\omega du dv$  for some real-valued function  $\omega : M \longrightarrow \mathbb{R}$ . Let  $\eta = e^{\frac{\omega}{2}} du$  and  $\xi = e^{\frac{\omega}{2}} dv$ . Then one can choose a lifting  $g = (g_1, g_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  such that the associated frame field  $\{e_0(g)\}$  is adapted with  $g^*(\omega^+) = -\eta$  and

$g^*(\omega^-) = -\xi$ . Since  $g^*(\omega^3) = 0$ ,

$$\begin{aligned} g_1^{-1}dg_1 &= \frac{1}{2}g^*\begin{pmatrix} \omega_2^1 & \omega^+ - \pi^+ \\ \omega^- + \pi^- & -\omega_2^1 \end{pmatrix} \\ &= \frac{1}{2}g^*\begin{pmatrix} \omega_2^1 & -\omega^+ - \pi^+ \\ \omega^- + \pi^- & -\omega_2^1 \end{pmatrix} + \eta\begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}, \\ g_2^{-1}dg_2 &= \frac{1}{2}g^*\begin{pmatrix} -\omega_2^1 & \omega^- - \pi^- \\ \omega^+ + \pi^+ & \omega_2^1 \end{pmatrix} \\ &= \frac{1}{2}g^*\begin{pmatrix} -\omega_2^1 & -\omega^- - \pi^- \\ \omega^+ + \pi^+ & \omega_2^1 \end{pmatrix} + \xi\begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Let  $\zeta = \frac{1}{2}g^*\begin{pmatrix} \omega_2^1 & -\omega^+ - \pi^+ \\ \omega^- + \pi^- & -\omega_2^1 \end{pmatrix} \in \mathfrak{sl}_2\mathbb{R}$ . Then  $d\zeta = -\zeta \wedge \zeta$ . The equation  $d\zeta = -\zeta \wedge \zeta$  satisfies the integrability condition; hence, by the Frobenius Theorem, there exists a smooth map  $h : M \longrightarrow \mathrm{SL}_2\mathbb{R}$  such that  $\zeta = h^{-1}dh$ . Since  $h \in \mathrm{SL}_2\mathbb{R}$ , it can be written

$$h = \begin{pmatrix} p_1 & -q_2 \\ q_1 & p_2 \end{pmatrix}, \quad p_1, p_2, q_1, q_2 \in \mathbb{R}.$$

Set  $F_1 := g_1 h^{-1}$ . Then

$$\begin{aligned} F_1^{-1}dF_1 &= (g_1 h^{-1})^{-1}d(g_1 h^{-1}) \\ &= h(g_1^{-1}dg_1)h^{-1} + hdh^{-1} \\ &= h\left[\zeta + \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}\eta\right]h^{-1} + hdh^{-1} \\ &= h\begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}h^{-1}\eta \\ &= \begin{pmatrix} p_1q_1 & -p_1^2 \\ q_1^2 & -p_1q_1 \end{pmatrix}\eta. \end{aligned}$$

The differential  $d$  can be written as  $d = \partial' + \partial''$ , where  $\partial'$  is the Lorentz holomorphic part and  $\partial''$  is the Lorentz antiholomorphic part. Since  $\partial''F_1 = \frac{\partial F_1}{\partial v}dv = 0$ ,  $F_1$  is Lorentz holomorphic.

Set  $F_2 = g_2 h^t$ . Then

$$\begin{aligned}
F_2^{-1} dF_2 &= (g_2 h^t)^{-1} d(g_2 h^t) \\
&= (h^t)^{-1} (g_2^{-1} dg_2) h^t + (h^t)^{-1} dh^t \\
&= (h^t)^{-1} \left[ -\zeta^t + \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} \xi \right] h^t + (h^t)^{-1} dh^t \\
&= (h^t)^{-1} \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} h^t \xi \\
&= \begin{pmatrix} p_2 q_2 & -p_2^2 \\ q_2^2 & -p_2 q_2 \end{pmatrix} \xi.
\end{aligned}$$

Since  $\partial' F_2 = \frac{\partial F_2}{\partial u} du = 0$ ,  $F_2$  is Lorentz antiholomorphic. Finally,

$$F_1 F_2^t = g_1 h^{-1} (g_2 h^t)^t = g_1 g_2^t = \varphi.$$

□

*Remark 4.* Note that, in Theorem 6,  $\varphi = F_1 F_2^t$  has cmc 1 (cmc  $-1$ ) if the framing  $F$  is orientation preserving (orientation reversing). In order to prove Theorem 6 for orientation reversing framing  $F$ , one needs to take

$$\begin{aligned}
F_1^{-1} dF_1 &= \begin{pmatrix} p_1 q_1 & p_1^2 \\ -q_1^2 & -p_1 q_1 \end{pmatrix} \eta, \quad F_2^{-1} dF_2 = \begin{pmatrix} p_2 q_2 & p_2^2 \\ -q_2^2 & -p_2 q_2 \end{pmatrix} \xi, \quad \text{and} \\
h &= \begin{pmatrix} q_2 & -p_1 \\ p_2 & q_1 \end{pmatrix} \in \text{SL}_2 \mathbb{R} \text{ in the proof.}
\end{aligned}$$

*Remark 5.*

$$\begin{aligned}
ds_\varphi^2 &= \varphi^*(ds^2) \\
&= \langle d\varphi, d\varphi \rangle \\
&= \langle d(F_1 F_2^t), d(F_1 F_2^t)^t \rangle \\
&= -\det\{F_1^{-1} dF_1 + (F_2^{-1} dF_2)^t\}.
\end{aligned}$$

So,  $\varphi$  does not assume degenerate points if and only if

$$\det\{F_1^{-1} dF_1 + (F_2^{-1} dF_2)^t\} \neq 0.$$

## 7. TIMELIKE SURFACES OF CONSTANT MEAN CURVATURE $\pm 1$ IN ADS 3-SPACE $\mathbb{H}_1^3(-1)$ VIA THE GROUP ACTION $\nu$

Let  $F : M \longrightarrow \text{SL}_2 \mathbb{R} \times \text{SL}_2 \mathbb{R}$  be a lift of  $\mathcal{F}$  to  $\text{SL}_2 \mathbb{R} \times \text{SL}_2 \mathbb{R}$  via the Lie group action  $\nu$ , i.e.,  $\nu(F)(\mathbf{1}, \mathbf{i}, \mathbf{j}', \mathbf{k}') = \mathcal{F}$ . Let  $\Omega := \Omega_1 \oplus \Omega_2 \in \mathfrak{sl}_2 \mathbb{R} \oplus \mathfrak{sl}_2 \mathbb{R}$ , where  $\Omega_1 = F_1^{-1} dF_1$ ,  $\Omega_2 = (dF_2^{-1}) F_2 \in \mathfrak{sl}_2 \mathbb{R}$ . The Gauß and Mainardi-Codazzi equations are equivalent to Maurer-Cartan equation

$$(54) \quad d\Omega + \Omega \wedge \Omega = 0,$$

which is the null curvature (integrability) condition of the Maurer-Cartan form  $\Omega$ . Let  $F^*$  denote the pull-back map  $F^* : (\mathfrak{sl}_2\mathbb{R})^* \oplus (\mathfrak{sl}_2\mathbb{R})^* \longrightarrow T^*M$ . The Maurer-Cartan forms  $\Omega_1 = F_1^{-1}dF_1$ ,  $\Omega_2 = (dF_2^{-1})F_2$  can be written as the following equations:

$$(55) \quad F_1^{-1}dF_1 = \frac{1}{2}F^* \begin{pmatrix} \omega^3 + \omega_2^1 & \omega^1 + \omega^2 - \omega_3^1 - \omega_3^2 \\ -\omega_1 + \omega_2 - \omega_3^1 + \omega_3^2 & -\omega^3 - \omega_2^1 \end{pmatrix},$$

$$(56) \quad (dF_2^{-1})F_2 = \frac{1}{2}F^* \begin{pmatrix} \omega^3 - \omega_2^1 & \omega^1 + \omega^2 + \omega_3^1 + \omega_3^2 \\ -\omega^1 + \omega^2 + \omega_3^1 - \omega_3^2 & -\omega^3 + \omega_2^1 \end{pmatrix}.$$

**Theorem 7** (A Bryant type representation formula for timelike cmc  $\pm 1$  Surfaces in  $\mathbb{H}_1^3(-1)$ ). *Let  $M$  be an open 2-manifold and  $F = (F_1, F_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  an immersion such that*

- (1)  $F_1$  is Lorentz holomorphic, i.e.,  $(F_1)_v = 0$  and  $F_2$  is Lorentz antiholomorphic, i.e.,  $(F_2)_u = 0$ ,
- (2)  $\det F_1^{-1}dF_1 = \det(dF_2^{-1})F_2 = 0$ .

Then

$$(57) \quad \psi := \nu(F)(\mathbf{1}) = F_1 F_2^{-1}$$

is a smooth conformal timelike immersion into  $\mathbb{H}_1^3(-1)$  with cmc  $\pm 1$ . Conversely, let  $M$  be an oriented and simply-connected Lorentzian 2-manifold with globally defined null coordinates. If  $\psi : M \longrightarrow \mathbb{H}_1^3(-1)$  is a smooth conformal timelike immersion with cmc  $\pm 1$ , then there exists an immersion  $F = (F_1, F_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  such that  $F_1, F_2$  satisfy the conditions (1), (2), and  $\psi = F_1 F_2^{-1}$ .

*Proof.* We use the same  $\omega^+, \omega^-, \pi^+, \pi^-$  and  $\alpha_i, \beta_i, \gamma_i$ ,  $i = 1, 2$  as defined in the proof of Theorem 6. Then

$$F_1^{-1}dF_1 = \begin{pmatrix} \alpha_1 & \beta_1 \\ \gamma_1 & -\alpha_1 \end{pmatrix} \text{ and } (dF_2^{-1})F_2 = \begin{pmatrix} \alpha_2 & \gamma_2 \\ \beta_2 & -\alpha_2 \end{pmatrix}.$$

Here,

$$ds_\psi^2 = F^*(\omega^3)^2 + F^*(\omega^+)F^*(\omega^-) = 2\alpha_1\alpha_2 + \beta_1\beta_2 + \gamma_1\gamma_2$$

defines an induced metric of  $\psi$  since  $F$  is an immersion.

Let  $U \subset M$  be a simply-connected open set in which there exists a null coordinate system  $(u, v)$  such that  $ds_\psi^2 = e^\omega du dv$ , for some real-valued function  $\omega : U \longrightarrow \mathbb{R}$ .  $M$  is covered by such open sets. Let  $\eta = e^{\frac{\omega}{2}} du$  and  $\xi = e^{\frac{\omega}{2}} dv$ . Then, by exactly the same argument in the proof of Theorem 6, there exists a smooth map  $h : U \longrightarrow \mathrm{SL}_2\mathbb{R}$  given by  $h = \begin{pmatrix} p_1 & -q_2 \\ q_1 & p_2 \end{pmatrix}$  and

$$e_0 \circ Fh = e_0(F_1h, F_2h) = F_1h(F_2h)^{-1} = F_1F_2^{-1} = e_0 \circ F.$$

$$\begin{aligned}
(F_1 h)^{-1} d(F_1 h) &= h^{-1} (F_1^{-1} dF_1) h + h^{-1} dh \\
&= \begin{pmatrix} p_2 dp_1 + q_2 dq_1 & -p_2 dq_2 + q_2 dp_2 - \eta \\ -q_1 dp_1 + p_1 dq_1 & q_1 dq_2 + p_1 dp_2 \end{pmatrix}
\end{aligned}$$

and

$$\begin{aligned}
d(F_2 h)^{-1} F_2 h &= (dh^{-1})h + h^{-1} [(dF_2^{-1})F_2]h \\
&= \begin{pmatrix} p_1 dp_2 + q_1 dq_2 & -q_2 dp_2 + p_2 dq_2 \\ -p_1 dq_1 + q_1 dp_1 - \xi & q_2 dq_1 + p_2 dp_1 \end{pmatrix}.
\end{aligned}$$

It then follows that  $(Fh)^*(\omega^3) = 0$ ,  $(Fh)^*(\omega^+) = -\eta$  and  $(Fh)^*(\omega^-) = -\xi$ . Hence,  $Fh : U \rightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  is an oriented framing in  $U$  along the immersion  $\psi = F_1 F_2^{-1}$  and  $\psi$  satisfies  $H = 1$  in  $U$ .

Conversely, let  $M$  be an oriented and simply-connected Lorentzian 2-manifold with globally defined null coordinates. Let  $\psi : M \rightarrow \mathbb{H}_1^3(-1)$  be a smooth conformal timelike immersion with cmc 1. There exist a null coordinate system  $(u, v)$  in  $M$  such that  $ds_\psi^2 = e^\omega du dv$  for some real-valued function  $\omega : M \rightarrow \mathbb{R}$ . Let  $\eta = e^{\frac{\omega}{2}} du$  and  $\xi = e^{\frac{\omega}{2}} dv$ . Then one can choose a lifting  $g : M \rightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  such that the associated frame field  $\{e_0(g)\}$  is adapted with  $g^*(\omega^+) = -\eta$  and  $g^*(\omega^-) = -\xi$ . Since  $g^*(\omega^3) = 0$ ,

$$\begin{aligned}
g_1^{-1} dg_1 &= \frac{1}{2} g^* \begin{pmatrix} \omega_2^1 & \omega^+ - \pi^+ \\ \omega^- + \pi^- & -\omega_2^1 \end{pmatrix} \\
&= \frac{1}{2} g^* \begin{pmatrix} \omega_2^1 & -\omega^+ - \pi^+ \\ \omega^- + \pi^- & -\omega_2^1 \end{pmatrix} + \eta \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}
\end{aligned}$$

and

$$\begin{aligned}
(dg_2^{-1})g_2 &= \frac{1}{2} g^* \begin{pmatrix} -\omega_2^1 & \omega^+ + \pi^+ \\ \omega^- - \pi^- & \omega_2^1 \end{pmatrix} \\
&= \frac{1}{2} g^* \begin{pmatrix} -\omega_2^1 & \omega^+ + \pi^+ \\ -\omega^- - \pi^- & \omega_2^1 \end{pmatrix} + \xi \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}.
\end{aligned}$$

Let  $\zeta = \frac{1}{2} g^* \begin{pmatrix} \omega_2^1 & -\omega^+ - \pi^+ \\ \omega^- + \pi^- & -\omega_2^1 \end{pmatrix} \in \mathfrak{sl}_2\mathbb{R}$ . Then  $d\zeta = -\zeta \wedge \zeta$  and this equation satisfies the integrability condition; hence, by the Frobenius Theorem, there exists a smooth map  $h : M \rightarrow \mathrm{SL}_2\mathbb{R}$  such that  $\zeta = h^{-1} dh$ . Since  $h \in \mathrm{SL}_2\mathbb{R}$ , it can be written  $h = \begin{pmatrix} p_1 & -q_2 \\ q_1 & p_2 \end{pmatrix}$ ,  $p_1, p_2, q_1, q_2 \in \mathbb{R}$ . Set  $F_1 = g_1 h^{-1}$ . Then

$$F_1^{-1} dF_1 = \begin{pmatrix} p_1 q_1 & -p_1^2 \\ q_1^2 & -p_1 q_1 \end{pmatrix} \eta$$

and  $F_1$  is Lorentz holomorphic. Set  $F_2 = g_2 h^{-1}$ . Then

$$\begin{aligned}
 (dF_2^{-1})F_2 &= (dh)h^{-1} + h[(dg_2^{-1})g_2]h^{-1} \\
 &= (dh)h^{-1} + h \left[ -\zeta + \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} \xi \right] h^{-1} \\
 &= h \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} h^{-1} \xi \\
 &= \begin{pmatrix} p_2 q_2 & q_2^2 \\ -p_2^2 & -p_2 q_2 \end{pmatrix} \xi
 \end{aligned}$$

and  $F_2$  is Lorentz antiholomorphic. Finally,

$$F_1 F_2^{-1} = g_1 h^{-1} (g_2 h^{-1})^{-1} = g_1 g_2^{-1} = \psi.$$

□

*Remark 6.* Note that

$$\begin{aligned}
 ds_\psi^2 &= \langle d\psi, d\psi \rangle \\
 &= \langle d(F_1 F_2^{-1}), d(F_1 F_2^{-1}) \rangle \\
 &= -\det\{F_1^{-1} dF_1 + (dF_2^{-1})F_2\}.
 \end{aligned}$$

So,  $\psi$  does not assume degenerate points if and only if

$$\det\{F_1^{-1} dF_1 + (dF_2^{-1})F_2\} \neq 0.$$

## 8. TIMELIKE MINIMAL SURFACES IN $\mathbb{E}_1^3$ AND THE CLASSICAL GAUSS MAP

Recall that the Lie group  $G \cong \mathrm{SL}_2\mathbb{R}$  acts isometrically on Lorentz 3-space  $\mathbb{E}_1^3$  via the Ad-action. The  $\mathrm{Ad}(G)$ -orbit of  $\mathbf{k}' = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  is a pseudosphere or de Sitter 2-space:

$$\mathbb{S}_1^2(1) = \{(x_1, x_2, x_3) \in \mathbb{E}_1^3 : -x_1^2 + x_2^2 + x_3^2 = 1\}.$$

The Ad-action of  $G$  on  $\mathbb{S}_1^2(1)$  is transitive as well. The isotropy subgroup of  $G$  at  $\mathbf{k}'$  is the indefinite orthogonal group  $\mathrm{SO}_1(2) = \{x_0 \mathbf{1} + x_3 \mathbf{k}' : x_0^2 - x_3^2 = 1\}$ . Thus,  $\mathbb{S}_1^2(1)$  can be identified with the symmetric space

$$\mathrm{SL}_2\mathbb{R}/\mathrm{SO}_1(2) = \left\{ h \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} h^{-1} : h \in \mathrm{SL}_2\mathbb{R} \right\}.$$

The orthogonal subgroup  $\mathrm{SO}_1(2)$  can be regarded as the hyperbola  $H_0^1$  in a Lorentz plane  $\mathbb{E}_1^2(x_0, x_3)$ . (This is a Lorentz analogue of  $S^1 \subset \mathbb{E}^2$ .) Note that the group  $H_0^1$  is isomorphic to the multiplicative group  $\mathbb{R}^\times = (\mathbb{R} \setminus \{0\}, \times)$ .

Let  $N = (0, 0, 1)$  and  $S = (0, 0, -1) \in \mathbb{S}_1^2(1)$  be the *north* and *south* pole of  $\mathbb{S}_1^2(1)$ . Let  $\wp_+ : \mathbb{S}_1^2(1) \setminus \{x_3 = -1\} \longrightarrow \mathbb{E}_1^2 \setminus H_0^1$  be the stereographic projection

from the south pole  $S = (0, 0, -1)$ , where  $H_0^1 = \{(x_1, x_2) \in \mathbb{E}_1^2 : -x_1^2 + x_2^2 = -1\}$ . Then

$$\begin{aligned}\wp_+(x_1, x_2, x_3) &= \left( \frac{x_1}{1+x_3}, \frac{x_2}{1+x_3} \right) \\ &\cong \left( \frac{x_1+x_2}{1+x_3}, \frac{-x_1+x_2}{1+x_3} \right) \in \mathbb{E}_1^2(u, v).\end{aligned}$$

Let  $\wp_- : \mathbb{S}_1^2(1) \setminus \{x_3 = 1\} \longrightarrow \mathbb{E}_1^2 \setminus H_0^1$  be the stereographic projection from the north pole  $S = (0, 0, 1)$ . Then

$$\begin{aligned}\wp_-(x_1, x_2, x_3) &= \left( \frac{x_1}{1-x_3}, \frac{x_2}{1-x_3} \right) \\ &\cong \left( \frac{x_1+x_2}{1-x_3}, \frac{-x_1+x_2}{1-x_3} \right) \in \mathbb{E}_1^2(u, v).\end{aligned}$$

Note that the classical Gauß map (i.e., the unit normal vector field)  $N$  of time-like surfaces in  $\mathbb{E}_1^3$  is mapped into de Sitter 2-space  $\mathbb{S}_1^2(1)$ . Thus, the image of classical Gauß map can be represented by the matrices  $h \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} h^{-1}$ ,  $h \in \text{SL}_2\mathbb{R}$ . If the timelike surface preserves the orientation, then  $h = \begin{pmatrix} p_1 & -q_2 \\ q_1 & p_2 \end{pmatrix} \in \text{SL}_2\mathbb{R}$ . So,

$$h \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} h^{-1} = \begin{pmatrix} p_1 p_2 - q_1 q_2 & 2p_1 q_2 \\ 2p_2 q_1 & -p_1 p_2 + q_1 q_2 \end{pmatrix} \in \mathbb{S}_1^2(1)$$

and  $\wp_-(h \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} h^{-1}) = \left( \frac{p_1}{q_1}, \frac{p_2}{q_2} \right) \in \mathbb{E}_1^2(u, v)$ . If the timelike surface reverses the orientation, then  $h = \begin{pmatrix} q_2 & -p_1 \\ p_2 & q_1 \end{pmatrix} \in \text{SL}_2\mathbb{R}$ . So,

$$h \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} h^{-1} = \begin{pmatrix} -p_1 p_2 + q_1 q_2 & 2p_1 q_2 \\ 2p_2 q_1 & p_1 p_2 - q_1 q_2 \end{pmatrix} \in \mathbb{S}_1^2(1)$$

and  $\wp_+(h \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} h^{-1}) = \left( \frac{p_1}{q_1}, \frac{p_2}{q_2} \right) \in \mathbb{E}_1^2(u, v)$ .

Here, we recall the following *Weierstraß formula* for a timelike minimal surface  $\psi : M \longrightarrow \mathbb{E}_1^3$  with data  $(q, f(q))$  and  $(r, g(v))$ :

$$(58) \quad \psi_u = \left( \frac{1}{2}(1+q^2), -\frac{1}{2}(1-q^2), -q \right) f(u), \quad \psi_v = \left( -\frac{1}{2}(1+r^2), -\frac{1}{2}(1-r^2), -r \right) g(v).$$

The induced metric of  $\psi$  is

$$ds_\psi^2 = (1+qr)^2 f(u)g(v)dudv.$$

*Remark 7.* The ordered pair  $(q, r)$  coincides with the projected Gauß map  $\wp_- \circ N$  of a timelike minimal surface with data  $(q, f(u))$  and  $(r, g(v))$ .

*Remark 8.* In [13], J. Inoguchi and M. Toda studied the construction of time-like minimal surfaces via loop group method. Their *normalized Wierstraß formula* for a timelike minimal surface  $\psi : M \longrightarrow \mathbb{E}_1^3$  with data  $(q, r)$  is

$$(59) \quad \psi_u = \left(\frac{1}{2}(1+q^2), -\frac{1}{2}(1-q^2), -q\right), \quad \psi_v = \left(-\frac{1}{2}(1+r^2), -\frac{1}{2}(1-r^2), -r\right)$$

and the induced metric of  $\psi$  is

$$ds_\psi^2 = (1+qr)^2 dudv.$$

In [13], the signs of coordinate functions in  $\psi_u$  and  $\psi_v$  are different. The reason is, in [13],  $(x_1, x_2, x_3) \in \mathbb{E}_1^3$  is identified with the matrix

$$\begin{pmatrix} -x_3 & -x_1+x_2 \\ x_1+x_2 & x_3 \end{pmatrix}, \text{ while in this paper it is identified with } \begin{pmatrix} x_3 & x_1+x_2 \\ -x_1+x_2 & -x_3 \end{pmatrix}.$$

Originally, this formula was obtained by M. A. Magid in [17]. However, in [17], the geometric meaning of the data  $(q, r)$  is not clarified. In [13], the data  $(q, r)$  are retrieved from the *normalized potential* in their construction. Moreover,  $q$  and  $r$  are the primitive functions of the coefficients  $Q$  and  $R$ , resp., of Hopf pairs. Note that this is locally true. In general,

$$(60) \quad q_u = \frac{Q}{f(u)},$$

$$(61) \quad r_v = \frac{R}{g(v)}.$$

As is mentioned in Remark 7,  $(q, r)$  is the projected Gauß map  $\wp_- \circ N$  of a timelike minimal surface given by the Weierstraß formula (59).

#### 9. LAWSON-GUICHARD CORRESPONDENCE BETWEEN TIMELIKE **cmc** $\pm 1$ SURFACES IN $\mathbb{H}_1^3(-1)$ AND TIMELIKE MINIMAL SURFACES IN $\mathbb{E}_1^3$

In Section 5, we discussed the Lawson-Guichard correspondence between timelike **cmc** surfaces in three different semi-Riemannian space forms  $\mathbb{E}_1^3$ ,  $\mathbb{S}_1^3(1)$  and  $\mathbb{H}_1^3(-1)$ . In particular, there is a one-to-one correspondence between timelike **cmc**  $\pm 1$  surfaces in  $\mathbb{H}_1^3(-1)$  and timelike minimal surfaces in  $\mathbb{E}_1^3$ . In this section, we give such bijective correspondence explicitly using the Bryant type representation formulae in Sections 6 and 7.

Let  $\varphi : M \longrightarrow \mathbb{H}^3(-1)$  be a timelike **cmc**  $-1$  surface. Then, by Theorem 6 (or by Theorem 7), there exists a smooth immersion  $F = (F_1, F_2) : M \longrightarrow \text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  such that

- (1)  $F_1$  is Lorentz holomorphic and  $F_2$  is Lorentz antiholomorphic.
- (2)  $\det(F_1^{-1}dF_1) = \det(F_2^{-1}dF_2) = 0$   
(or (2')  $\det(F_1^{-1}dF_1) = \det((dF_2^{-1})F_2) = 0$ ).
- (3)  $\varphi = F_1 F_2^t$  (or (3')  $\varphi = F_1 F_2^{-1}$ ).

As we have seen in the proof of Theorem 6 (or Theorem 7), locally in an open set  $U \subset M$ ,

$$\begin{aligned} F_1^{-1}dF_1 &= \begin{pmatrix} p_1 q_1 & -p_1^2 \\ q_1^2 & -p_1 q_1 \end{pmatrix} \eta \\ &= \begin{pmatrix} \frac{p_1}{q_1} & -\frac{p_1^2}{q_1^2} \\ 1 & -\frac{p_1}{q_1} \end{pmatrix} q_1^2 \eta \end{aligned}$$

and similarly,

$$F_2^{-1}dF_2 = \begin{pmatrix} \frac{p_2}{q_2} & -\frac{p_2^2}{q_2^2} \\ 1 & -\frac{p_2}{q_2} \end{pmatrix} q_2^2 \xi$$

or

$$(dF_2^{-1})F_2 = \begin{pmatrix} \frac{p_2}{q_2} & 1 \\ -\frac{p_2^2}{q_2^2} & -\frac{p_2}{q_2} \end{pmatrix} q_2^2 \xi.$$

Let  $q := \frac{p_1}{q_1}$ ,  $f(u) := q_1^2$ ,  $r := \frac{p_2}{q_2}$ , and  $g(v) := q_2^2$ . Then the Weierstraß formula (58) defines a timelike minimal surface  $\psi : U \longrightarrow \mathbb{E}_1^3$ .

$$\begin{aligned} ds_\varphi^2 &= \eta \xi \\ &= e^\omega dudv \\ &= e^\omega (1 + qr)^2 f(u)g(v)dudv. \end{aligned}$$

So, the induced metric  $ds_\varphi^2$  of the timelike cmc  $-1$  surface  $\varphi$  is conformal to  $ds_\psi^2$ .

Conversely, assume that a timelike minimal surface  $\psi : M \longrightarrow \mathbb{E}_1^3$  is given by the Weierstraß formula (58) with data  $(q, f(u))$  and  $(r, g(v))$ . Consider the following system of differential equations:

$$(62) \quad F_1^{-1}dF_1 = \begin{pmatrix} q & -q^2 \\ 1 & -q \end{pmatrix} f(u)du, \quad F_2^{-1}dF_2 = \begin{pmatrix} r & -r^2 \\ 1 & -r \end{pmatrix} g(v)dv.$$

Since these equations satisfy the integrability condition, there exists a solution  $(F_1, F_2) : M \longrightarrow \text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  satisfying the conditions (1) and (2). By Theorem 6,  $\varphi := F_1 F_2^t : M \longrightarrow \mathbb{H}_1^3(-1)$  defines a timelike cmc  $-1$  surface in  $\mathbb{H}_1^3(-1)$ . Similarly, the system of differential equations:

$$(63) \quad F_1^{-1}dF_1 = \begin{pmatrix} q & -q^2 \\ 1 & -q \end{pmatrix} f(u)du, \quad (dF_2^{-1})F_2 = \begin{pmatrix} r & 1 \\ -r^2 & -r \end{pmatrix} g(v)dv$$

satisfies the integrability condition; hence there exists a solution  $(F_1, F_2) : M \longrightarrow \text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  satisfying conditions (1) and (2'). By Theorem 7,  $\varphi := F_1 F_2^{-1}$  defines a timelike cmc  $-1$  surface in  $\mathbb{H}_1^3(-1)$ .

### 10. THE HYPERBOLIC GAUSS MAP OF TIMELIKE SURFACES IN $\mathbb{H}_1^3(-1)$

Let  $\varphi : M \rightarrow \mathbb{H}_1^3(-1)$  be an oriented timelike surface in  $\mathbb{H}_1^3(-1)$ . At each base point  $e_0 = \varphi(m) \in \mathbb{H}_1^3(-1)$ ,  $e_3 \in T_{e_0}\mathbb{H}_1^3(-1)$  is an oriented unit normal vector to the tangent plane  $\varphi_*(T_m M)$ . The oriented normal geodesic in  $\mathbb{H}_1^3(-1)$  emanating from  $e_0$ , which is tangent to the normal vector  $e_3(\varphi(m))$  asymptotically approaches to the *null cone*  $\mathbb{N} = \{\mathbf{u} \in \mathbb{E}_2^4 : \langle \mathbf{u}, \mathbf{u} \rangle = 0\}$  at exactly two points  $[e_0 + e_3], [e_0 - e_3] \in \mathbb{N}$ . The orientation allows us to name  $[e_0 + e_3]$  the initial point and  $[e_0 - e_3]$  the terminal point.

Define a map  $G : M \rightarrow \mathbb{N}$  by  $G(m) = [e_0 + e_3](m)$  for each  $m \in M$ . This map is an analogue of the hyperbolic Gauß map<sup>7</sup> of surfaces in hyperbolic 3-space  $\mathbb{H}_1^3(-1)$ . The map will still be called the *hyperbolic Gauß map* here.

Let  $\varphi : M \rightarrow \mathbb{H}_1^3(-1)$  be a timelike surface in  $\mathbb{H}_1^3(-1)$  and  $d\sigma^2$  denote the induced metric on  $\mathbb{N}^3$ . Then

$$d\sigma_\varphi^2 := (e_0 + e_3)^*(d\sigma^2) = \langle d(e_0 + e_3), d(e_0 + e_3) \rangle = -(\omega^1 + \omega_3^1)^2 + (\omega^2 + \omega_3^2)^2.$$

**Proposition 8.** *The hyperbolic Gauß map  $[e_0 + e_3] : M \rightarrow \mathbb{N}$  ( $[e_0 - e_3] : M \rightarrow \mathbb{N}$ ) of a timelike surface  $\varphi : M \rightarrow \mathbb{H}_1^3(-1)$  is conformal if and only if  $\varphi$  satisfies  $H = 1$  ( $H = -1$ ) or  $\varphi$  is totally umbilic.*

*Proof.* We will assume the same settings in the proof of Proposition 1. Then

$$\begin{aligned} d\sigma_\varphi^2 &= -(\omega^1 + \omega_3^1)^2 + (\omega^2 + \omega_3^2)^2 \\ &= -((1 - h_{11})\omega^1 + h_{12}\omega^2)^2 + (-h_{12}\omega^1 + (1 - h_{22})\omega^2)^2 \\ &= (2H(H - 1) - K)ds_\varphi^2 - \\ &\quad (H - 1)((h_{11} - h_{22})(\omega^1)^2 - 4h_{12}\omega^1 \otimes \omega^2 + (h_{11} - h_{22})(\omega^2)^2). \end{aligned}$$

Thus,  $[e_0 + e_3]$  is conformal, i.e.,  $d\sigma_\varphi^2$  is a multiple of  $ds_\varphi^2$  if and only if  $(H - 1)(h_{11} - h_{22}) = (H - 1)h_{12} = 0$ .

If  $H = 1$ , then  $d\sigma_\varphi^2 = -Kds_\varphi^2$ . Suppose  $H \neq 1$ . Let  $U = \{m \in M : H(m) \neq 1\}$ . Then  $U$  is clearly open in  $M$ . Since  $H \neq 1$  on  $U$ ,  $h_{11} = h_{22}$  and  $h_{12} = 0$  on  $U$ . The second fundamental form  $II$  is then

$$\begin{aligned} II &= -h_{11}(\omega^1)^2 + 2h_{12}\omega^1 \otimes \omega^2 + h_{22}(\omega^2)^2 \\ &= h_{11}(-(\omega^1)^2 + (\omega^2)^2) \\ &= HI \text{ on } U. \end{aligned}$$

By comparing with the equation (1), we see that the Hopf differential  $\mathcal{Q} = 0$  on  $U$ , i.e.,  $\varphi(U)$  is totally umbilic. Note that  $H$  must be constant on  $U$ , since  $Q = R = 0$ . Let  $V$  be a connected component of  $U$ . Since  $H$  is constant on  $V$  and  $H$  is continuous on  $M$ ,  $H$  is constant on  $\bar{V}$ . This implies that  $H \neq 1$  on  $\bar{V}$  and so  $\bar{V} \subset U$ . Since  $V$  is connected, so is  $\bar{V}$ . However,  $V$  is a connected

---

<sup>7</sup>The hyperbolic Gauß map was introduced by C. Epstein ([7]) and was used by R. L. Bryant in his study of cmc 1 surfaces in hyperbolic 3-space  $\mathbb{H}^3(-1)$  ([4]).

component; thus,  $V = \bar{V}$ . It then follows from the connectedness of  $M$  that  $M = V$ . Therefore,  $\varphi$  is totally umbilic on  $M$ . The converse is trivial.  $\square$

*Remark 9.* Note that  $[e_0 + e_3] : M \rightarrow \mathbb{N}$  ( $[e_0 - e_3] : M \rightarrow \mathbb{N}$ ) is the hyperbolic Gauß map of a timelike surface  $\varphi : M \rightarrow \mathbb{H}_1^3(-1)$  if  $\varphi$  preserves (reverses) the orientation.

*Remark 10.* If  $\varphi$  satisfies  $\pm 1$  and is totally umbilic, then the hyperbolic Gauß map is constant, since  $d\sigma_\varphi^2 = 0$ . It is also shown in Section 13 that if  $\varphi$  satisfies  $\pm 1$  and is totally umbilic, then the (projected) hyperbolic Gauß map is both Lorentz holomorphic and antiholomorphic; hence it is constant. By the equation (2), the Gaussian curvature  $K = H^2 - 1 = 0$ . Thus,  $\varphi$  may be regarded as a *horosphere* type surface in  $\mathbb{H}_1^3(-1)$ .

The null cone  $\mathbb{N}$  satisfies the quadric equation

$$-(x_0)^2 - (x_1)^2 + (x_2)^2 + (x_3)^2 = 0.$$

If  $x_0 \neq 0$ , then the above equation can be written

$$-\left(\frac{x_1}{x_0}\right)^2 + \left(\frac{x_2}{x_0}\right)^2 + \left(\frac{x_3}{x_0}\right)^2 = 1,$$

i.e.,  $\mathbb{N}$  can be locally identified with de Sitter 2-space  $\mathbb{S}_1^2(1)$ . With this identification, the hyperbolic Gauß map can be mapped into de Sitter 2-space  $\mathbb{S}_1^2(1)$ . Thus, we may be able to relate the hyperbolic Gauß map of a timelike cmc  $\pm 1$  surface in  $\mathbb{H}_1^3(-1)$  and the Gauß map of corresponding timelike minimal surface in  $\mathbb{E}_1^3$ . This relationship is discussed in the next section.

Let  $(x_0, x_1, x_2, x_3) \in \mathbb{N}$ . Denote by  $[x_0, x_1, x_2, x_3]$  the null line generated by the null vector  $(x_0, x_1, x_2, x_3)$ . By using nonhomogeneous coordinates,

$$\begin{aligned} [x_0, x_1, x_2, x_3] &= \left[1, \frac{x_1}{x_0}, \frac{x_2}{x_0}, \frac{x_3}{x_0}\right] \text{ provided } x_0 \neq 0 \\ &\cong \left(\frac{x_1}{x_0}, \frac{x_2}{x_0}, \frac{x_3}{x_0}\right) \in \mathbb{S}_1^2(1) \\ &\cong \left(\frac{x_1}{x_0 + x_3}, \frac{x_2}{x_0 + x_3}\right) \in \mathbb{E}_1^2 \text{ via the projection } \wp_+ \end{aligned}$$

or

$$[x_0, x_1, x_2, x_3] \cong \left(\frac{x_1}{x_0 - x_3}, \frac{x_2}{x_0 - x_3}\right) \in \mathbb{E}_1^2 \text{ via the projection } \wp_-.$$

Finally, we have the identification:

$$(64) \quad [x_0, x_1, x_2, x_3] \cong \left(\frac{x_1 + x_2}{x_0 + x_3}, \frac{-x_1 + x_2}{x_0 + x_3}\right) \in \mathbb{E}_1^2(u, v)$$

or

$$(65) \quad [x_0, x_1, x_2, x_3] \cong \left(\frac{x_1 + x_2}{x_0 - x_3}, \frac{-x_1 + x_2}{x_0 - x_3}\right) \in \mathbb{E}_1^2(u, v).$$

## 11. THE HYPERBOLIC GAUSS MAP AND THE SECONDARY GAUSS MAP

Let  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  be a timelike **cmc**  $-1$  surface in  $\mathbb{H}_1^3(-1)$ . Then by Theorem 6 (or by Theorem 7), there exists a smooth immersion  $F = (F_1, F_2) : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  such that

- (1)  $F_1$  is Lorentz holomorphic and  $F_2$  is Lorentz antiholomorphic.
- (2)  $\det(F_1^{-1}dF_1) = \det(F_2^{-1}dF_2) = 0$   
(or (2')  $\det(F_1^{-1}dF_1) = \det((dF_2^{-1})F_2) = 0$ ).
- (3)  $\varphi = F_1 F_2^t$  (or (3')  $\varphi = F_1 F_2^{-1}$ ).

Let  $F_1 = \begin{pmatrix} F_{11} & F_{12} \\ F_{13} & F_{14} \end{pmatrix}$  and  $F_2 = \begin{pmatrix} F_{21} & F_{22} \\ F_{23} & F_{24} \end{pmatrix}$ . Then

$$\begin{aligned} (e_0 + e_3)(F) &= F_1(\mathbf{1} + \mathbf{k}')F_2^t \\ &= 2 \begin{pmatrix} F_{11} & F_{12} \\ F_{13} & F_{14} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} F_{21} & F_{23} \\ F_{22} & F_{24} \end{pmatrix} \\ &= 2 \begin{pmatrix} F_{11}F_{21} & F_{11}F_{23} \\ F_{13}F_{21} & F_{13}F_{23} \end{pmatrix}. \end{aligned}$$

By the identification (65),

$$\begin{aligned} [(e_0 + e_3)(F)] &= \left[ \begin{pmatrix} F_{11}F_{21} & F_{11}F_{23} \\ F_{13}F_{21} & F_{13}F_{23} \end{pmatrix} \right] \\ &\cong \left( \frac{F_{11}F_{23}}{F_{13}F_{23}}, \frac{F_{13}F_{21}}{F_{13}F_{23}} \right) \\ &= \left( \frac{F_{11}}{F_{13}}, \frac{F_{21}}{F_{23}} \right) \in \mathbb{E}_1^2(u, v). \end{aligned}$$

Similarly,

$$\begin{aligned} [(e_0 - e_3)(F)] &= [F_1(\mathbf{1} - \mathbf{k}')F_2^t] \\ &\cong \left( \frac{F_{12}}{F_{14}}, \frac{F_{22}}{F_{24}} \right) \in \mathbb{E}_1^2(u, v). \end{aligned}$$

If  $(e_0 \pm e_3)(F) = F_1(\mathbf{1} \pm \mathbf{k}')F_2^{-1}$ , then

$$\begin{aligned} [(e_0 + e_3)(F)] &\cong \left( \frac{F_{11}}{F_{13}}, -\frac{F_{24}}{F_{22}} \right) \in \mathbb{E}_1^2(u, v), \\ [(e_0 - e_3)(F)] &\cong \left( \frac{F_{12}}{F_{14}}, -\frac{F_{23}}{F_{21}} \right) \in \mathbb{E}_1^2(u, v). \end{aligned}$$

Locally,

$$\begin{aligned} (e_0 + e_3)(Fh) &= F_1 h(\mathbf{1} + \mathbf{k}') (F_2 (h^{-1})^t)^t \\ &= 2F_1 h \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} h^{-1} F_2^t \\ &= 2F_1 \begin{pmatrix} p_1 & -q_2 \\ q_1 & p_2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} p_2 & q_2 \\ -q_1 & p_1 \end{pmatrix} \begin{pmatrix} F_{21} & F_{23} \\ F_{22} & F_{24} \end{pmatrix}. \end{aligned}$$

The last expression is simplified to the matrix:

$$2 \begin{pmatrix} (F_{11}p_1 + F_{12}q_1)(F_{21}p_2 + F_{22}q_2) & (F_{11}p_1 + F_{12}q_1)(F_{23}p_2 + F_{24}q_2) \\ (F_{13}p_1 + F_{14}q_1)(F_{21}p_2 + F_{22}q_2) & (F_{13}p_1 + F_{14}q_1)(F_{23}p_2 + F_{24}q_2) \end{pmatrix}.$$

Thus, by the identification (65),

$$\begin{aligned} [(e_0 + e_3)(Fh)] &= [F_1 h(\mathbf{1} + \mathbf{k}') (F_2 (h^{-1})^t)^t] \\ &\cong \left( \frac{F_{11}p_1 + F_{12}q_1}{F_{13}p_1 + F_{14}q_1}, \frac{F_{21}p_2 + F_{22}q_2}{F_{23}p_2 + F_{24}q_2} \right) \in \mathbb{E}_1^2(u, v). \end{aligned}$$

Note that the hyperbolic Gauß map  $[e_0 + e_3]$  is orientation preserving while  $[e_0 - e_3]$  is orientation reversing. So,

$$\begin{aligned} [(e_0 - e_3)(Fh)] &= [F_1 h(\mathbf{1} - \mathbf{k}') (F_2 (h^{-1})^t)^t] \\ &\cong \left( \frac{F_{11}p_1 - F_{12}q_1}{F_{13}p_1 - F_{14}q_1}, \frac{F_{21}p_2 - F_{22}q_2}{F_{23}p_2 - F_{24}q_2} \right) \in \mathbb{E}_1^2(u, v), \end{aligned}$$

where  $h = \begin{pmatrix} q_2 & -p_1 \\ p_2 & q_1 \end{pmatrix}$ . (See Remark 4.)

Similarly, if  $(e_0 \pm e_3)(Fh) = F_1 h(\mathbf{1} \pm \mathbf{k}') (F_2 h)^{-1}$ , then

$$\begin{aligned} [(e_0 + e_3)(Fh)] &\cong \left( \frac{F_{11}p_1 + F_{12}q_1}{F_{13}p_1 + F_{14}q_1}, -\frac{F_{24}p_2 - F_{23}q_2}{F_{22}p_2 - F_{21}q_2} \right) \in \mathbb{E}_1^2(u, v), \\ [(e_0 - e_3)(Fh)] &\cong \left( \frac{F_{11}p_1 - F_{12}q_1}{F_{13}p_1 - F_{14}q_1}, -\frac{F_{24}p_2 + F_{23}q_2}{F_{22}p_2 + F_{21}q_2} \right) \in \mathbb{E}_1^2(u, v). \end{aligned}$$

Let  $q := \frac{p_1}{q_1}$  and  $r := \frac{p_2}{q_2}$ . The ordered pair  $(q, r)$  is called the *secondary Gauß map*<sup>8</sup>. In terms of the secondary Gauß map  $(q, r)$ , locally in an open

---

<sup>8</sup>As seen in section 8, this secondary Gauß map coincides with the projected Gauß map of a corresponding timelike minimal surface in  $\mathbb{E}_1^3$ .

set  $U \subset M$ ,

$$\begin{aligned} F_1^{-1}dF_1 &= \begin{pmatrix} q & -q^2 \\ 1 & -q \end{pmatrix} f(u)du = \begin{pmatrix} q & -q^2 \\ 1 & -q \end{pmatrix} \eta, \\ F_2^{-1}dF_2 &= \begin{pmatrix} r & -r^2 \\ 1 & -r \end{pmatrix} g(v)dv = \begin{pmatrix} r & -r^2 \\ 1 & -r \end{pmatrix} \xi, \\ (dF_2^{-1})F_2 &= \begin{pmatrix} r & 1 \\ -r^2 & -r \end{pmatrix} g(v)dv = \begin{pmatrix} r & 1 \\ -r^2 & -r \end{pmatrix} \xi. \end{aligned}$$

Thus, we have the following equations:

$$\begin{aligned} dF_1 &= \begin{pmatrix} dF_{11} & dF_{12} \\ dF_{13} & dF_{14} \end{pmatrix} = \begin{pmatrix} F_{11} & F_{12} \\ F_{13} & F_{14} \end{pmatrix} \begin{pmatrix} q & -q^2 \\ 1 & -q \end{pmatrix} \eta \\ &= \begin{pmatrix} F_{11}q + F_{12} & -(F_{11}q + F_{12}q) \\ F_{13}q + F_{14} & -(F_{13}q + F_{14}q) \end{pmatrix} \eta, \end{aligned}$$

$$\begin{aligned} dF_2 &= \begin{pmatrix} dF_{21} & dF_{22} \\ dF_{23} & dF_{24} \end{pmatrix} = \begin{pmatrix} F_{21} & F_{22} \\ F_{23} & F_{24} \end{pmatrix} \begin{pmatrix} r & -r^2 \\ 1 & -r \end{pmatrix} \xi \\ &= \begin{pmatrix} F_{21}r + F_{22} & -(F_{21}r + F_{22})r \\ F_{23}r + F_{24} & -(F_{23}r + F_{24})r \end{pmatrix} \xi \end{aligned}$$

and

$$\begin{aligned} dF_2^{-1} &= \begin{pmatrix} dF_{24} & -dF_{22} \\ -dF_{23} & dF_{21} \end{pmatrix} = \begin{pmatrix} r & 1 \\ -r^2 & -r \end{pmatrix} \begin{pmatrix} F_{24} & -F_{22} \\ -F_{23} & F_{21} \end{pmatrix} \xi \\ &= \begin{pmatrix} F_{24}r - F_{22} & -(F_{22}r - F_{21}) \\ -(F_{24}r - F_{23})r & (F_{22}r - F_{21})r \end{pmatrix} \xi, \end{aligned}$$

i.e.,  $dF_2$  is also given by

$$dF_2 = \begin{pmatrix} (F_{22}r - F_{21})r & F_{22}r - F_{21} \\ (F_{24}r - F_{23})r & F_{24}r - F_{23} \end{pmatrix} \xi.$$

Hence, the hyperbolic Gauß map can be written:

$$\begin{aligned}
[(e_0 + e_3)(Fh)] &= [F_1 h(\mathbf{1} + \mathbf{k}') (F_2 (h^{-1})^t)^t] \\
&\cong \left( \frac{F_{11}q + F_{12}}{F_{13}q + F_{14}}, \frac{F_{21}r + F_{22}}{F_{23}r + F_{24}} \right) \\
&= \left( \frac{dF_{11}}{dF_{13}}, \frac{dF_{21}}{dF_{23}} \right) \\
&= \left( \frac{dF_{12}}{dF_{14}}, \frac{dF_{22}}{dF_{24}} \right) \in \mathbb{E}_1^2(u, v), \\
[(e_0 + e_3)(Fh)] &= [F_1 h(\mathbf{1} + \mathbf{k}') (F_2 h)^{-1}] \\
&= \left( \frac{F_{11}q + F_{12}}{F_{13}q + F_{14}}, -\frac{F_{24}r - F_{23}}{F_{22}r - F_{21}} \right) \\
&= \left( \frac{dF_{11}}{dF_{13}}, -\frac{dF_{23}}{dF_{21}} \right) \\
&= \left( \frac{dF_{12}}{dF_{14}}, -\frac{dF_{24}}{dF_{22}} \right) \in \mathbb{E}_1^2(u, v)
\end{aligned}$$

and

$$\begin{aligned}
[(e_0 - e_3)(Fh)] &= [F_1 h(\mathbf{1} - \mathbf{k}') (F_2 (h^{-1})^t)^t] \\
&\cong \left( \frac{F_{11}q - F_{12}}{F_{13}q - F_{14}}, \frac{F_{21}r - F_{22}}{F_{23}r - F_{24}} \right) \\
&= \left( \frac{dF_{11}}{dF_{13}}, \frac{dF_{21}}{dF_{23}} \right) \\
&= \left( \frac{dF_{12}}{dF_{14}}, \frac{dF_{22}}{dF_{24}} \right) \in \mathbb{E}_1^2(u, v), \\
[(e_0 - e_3)(Fh)] &= [F_1 h(\mathbf{1} - \mathbf{k}') (F_2 h)^{-1}] \\
&\cong \left( \frac{F_{11}q - F_{12}}{F_{13}q - F_{14}}, -\frac{F_{24}r + F_{23}}{F_{22}r + F_{21}} \right) \\
&= \left( \frac{dF_{11}}{dF_{13}}, -\frac{dF_{23}}{dF_{21}} \right) \\
&= \left( \frac{dF_{12}}{dF_{14}}, -\frac{dF_{24}}{dF_{22}} \right) \in \mathbb{E}_1^2(u, v).
\end{aligned}$$

## 12. THE GENERALIZED GAUSS MAP AND THE HYPERBOLIC GAUSS MAP

Let  $G(2, \mathbb{E}_2^4)$  be the Grassmannian manifold of oriented timelike 2-planes in  $\mathbb{E}_2^4$ . Let  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  be an oriented timelike surface in  $\mathbb{H}_1^3(-1)$ . At each point  $p \in M$ , there is a (timelike) tangent plane to the surface  $\varphi$ :

$$\varphi_*(T_p M) = [\varphi_x \wedge \varphi_y]_p = [(\varphi_x + \varphi_y) \wedge (-\varphi_x + \varphi_y)]_p$$

spanned by a timelike vector  $\varphi_x$  and a spacelike vector  $\varphi_y$  or equivalently, by two null vectors  $\varphi_x + \varphi_y$  and  $-\varphi_x + \varphi_y$ . Define a map

$$\mathcal{G} : M \longrightarrow G(2, \mathbb{E}_2^4); p \in M \xrightarrow{\mathcal{G}} [\varphi_x \wedge \varphi_y]_p = [(\varphi_x + \varphi_y) \wedge (-\varphi_x + \varphi_y)]_p.$$

This map is called the *generalized Gauss map* of a timelike surface  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$ .

Let  $\mathbf{u}$  be a null vector, i.e.,  $\langle \mathbf{u}, \mathbf{u} \rangle = 0$  and  $[\mathbf{u}]$  denote the null line generated by  $\mathbf{u}$ . Let  $\mathbb{Q}_0^2 := \{[\mathbf{u}] \in \mathbb{RP}_2^3 : \langle \mathbf{u}, \mathbf{u} \rangle = 0\}$ . Then there is an embedding

$$\Gamma : G(2, \mathbb{E}_2^4) \longrightarrow \mathbb{Q}_0^2 \times \mathbb{Q}_0^2; [\mathbf{v} \wedge \mathbf{w}] \xrightarrow{\Gamma} ([\mathbf{v} + \mathbf{w}], [-\mathbf{v} + \mathbf{w}]),$$

where  $\mathbf{v}$  is a timelike vector and  $\mathbf{w}$  is a spacelike vector. That is,  $G(2, \mathbb{E}_2^4) \cong \text{Im} \Gamma \subset \mathbb{Q}_0^2 \times \mathbb{Q}_0^2$ . The Lie group  $\text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  acts on  $G(2, \mathbb{E}_2^4)$  transitively via the actions:

$$\begin{aligned} \mu : (\text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}) \times G(2, \mathbb{E}_2^4) &\longrightarrow G(2, \mathbb{E}_2^4); \\ \mu(g, [\mathbf{v} \wedge \mathbf{w}]) &:= [\mu(g, \mathbf{v}) \wedge \mu(g, \mathbf{w})] = [(g_1 \mathbf{v} g_2^t) \wedge (g_1 \mathbf{w} g_2^t)] \end{aligned}$$

and

$$\begin{aligned} \nu : (\text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}) \times G(2, \mathbb{E}_2^4) &\longrightarrow G(2, \mathbb{E}_2^4); \\ \nu(g, [\mathbf{v} \wedge \mathbf{w}]) &:= [\nu(g, \mathbf{v}) \wedge \nu(g, \mathbf{w})] = [(g_1 \mathbf{v} g_2^{-1}) \wedge (g_1 \mathbf{w} g_2^{-1})] \end{aligned}$$

for  $g = (g_1, g_2) \in \text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  and  $[\mathbf{v}, \mathbf{w}] \in G(2, \mathbb{E}_2^4)$ . Note that

$$\begin{aligned} \mu(g, [\mathbf{v} \wedge \mathbf{w}]) &\cong ([g_1(\mathbf{v} + \mathbf{w})g_2^t], [g_1(-\mathbf{v} + \mathbf{w})g_2^t]) \in \mathbb{Q}_0^2 \times \mathbb{Q}_0^2, \\ \nu(g, [\mathbf{v} \wedge \mathbf{w}]) &\cong ([g_1(\mathbf{v} + \mathbf{w})g_2^{-1}], [g_1(-\mathbf{v} + \mathbf{w})g_2^{-1}]) \in \mathbb{Q}_0^2 \times \mathbb{Q}_0^2. \end{aligned}$$

The isotropy subgroup of  $\text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  with the actions  $\mu$  and  $\nu$  at

$$\begin{aligned} [e_1 \wedge e_2] &= [(e_1 + e_2) \wedge (-e_1 + e_2)] \\ &= \left[ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \wedge \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right] \\ &\cong \left( \left[ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right], \left[ \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right] \right) \end{aligned}$$

is  $\mathbb{R}_+ \times \mathbb{R}_+$ , where  $\mathbb{R}_+ := \left\{ \begin{pmatrix} r & 0 \\ 0 & \frac{1}{r} \end{pmatrix} : r \in \mathbb{R} \setminus \{0\} \right\}$ . Thus, the Grassmannian manifold  $G(2, \mathbb{E}_2^4)$  can be represented as a symmetric space

$$\begin{aligned} &\frac{\text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}}{\mathbb{R}_+ \times \mathbb{R}_+} \cong \\ &\left\{ \left( \left[ g_1 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} g_2^t \right], \left[ g_1 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} g_2^t \right] \right) \in \mathbb{Q}_0^2 \times \mathbb{Q}_0^2 : g_1, g_2 \in \text{SL}_2\mathbb{R} \right\} \end{aligned}$$

or

$$\frac{\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}}{\mathbb{R}_+ \times \mathbb{R}_+} \cong \left\{ \left( \begin{bmatrix} 0 & 1 \\ g_1 & 0 \end{bmatrix} g_2^{-1}, \begin{bmatrix} 0 & 0 \\ g_1 & 1 \end{bmatrix} g_2^{-1} \right) \in \mathbb{Q}_0^2 \times \mathbb{Q}_0^2 : g_1, g_2 \in \mathrm{SL}_2\mathbb{R} \right\}.$$

Denote by  $G(2, \mathbb{E}_2^4)^-$  the Grassmannian manifold of *negatively* oriented time-like 2-planes in  $\mathbb{E}_2^4$ . Then

$$G(2, \mathbb{E}_2^4)^- = \frac{\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}}{\mathbb{R}_+ \times \mathbb{R}_+} \cong \left\{ \left( \begin{bmatrix} 0 & 0 \\ g_1 & 1 \end{bmatrix} g_2^t, \begin{bmatrix} 0 & 1 \\ g_1 & 0 \end{bmatrix} g_2^t \right) \in \mathbb{Q}_0^2 \times \mathbb{Q}_0^2 : g_1, g_2 \in \mathrm{SL}_2\mathbb{R} \right\}$$

or

$$G(2, \mathbb{E}_2^4)^- = \frac{\mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}}{\mathbb{R}_+ \times \mathbb{R}_+} \cong \left\{ \left( \begin{bmatrix} 0 & 0 \\ g_1 & 1 \end{bmatrix} g_2^{-1}, \begin{bmatrix} 0 & 1 \\ g_1 & 0 \end{bmatrix} g_2^{-1} \right) \in \mathbb{Q}_0^2 \times \mathbb{Q}_0^2 : g_1, g_2 \in \mathrm{SL}_2\mathbb{R} \right\}.$$

Let us write  $g_1 = \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix}$  and  $g_2 = \begin{pmatrix} g_{21} & g_{22} \\ g_{23} & g_{24} \end{pmatrix}$ . Define a projection map

$$\phi = (\phi_1, \phi_2) : G(2, \mathbb{E}_2^4) \longrightarrow \mathbb{E}_1^2(u, v) \times \mathbb{E}_1^2(u, v)$$

by

$$\left( \begin{bmatrix} 0 & 1 \\ g_1 & 0 \end{bmatrix} g_2^t, \begin{bmatrix} 0 & 0 \\ g_1 & 1 \end{bmatrix} g_2^t \right) \xrightarrow{(\phi_1, \phi_2)} \left( \frac{g_{11}}{g_{13}}, \frac{g_{21}}{g_{23}} \right)$$

or

$$\left( \begin{bmatrix} 0 & 1 \\ g_1 & 0 \end{bmatrix} g_2^{-1}, \begin{bmatrix} 0 & 0 \\ g_1 & 1 \end{bmatrix} g_2^{-1} \right) \xrightarrow{(\phi_1, \phi_2)} \left( \frac{g_{11}}{g_{13}}, -\frac{g_{24}}{g_{22}} \right).$$

Similarly, we also define a projection map

$$\phi^- = (\phi_1^-, \phi_2^-) : G(2, \mathbb{E}_2^4)^- \longrightarrow \mathbb{E}_1^2(u, v) \times \mathbb{E}_1^2(u, v)$$

by

$$\left( \begin{bmatrix} 0 & 0 \\ g_1 & 1 \end{bmatrix} g_2^t, \begin{bmatrix} 0 & 1 \\ g_1 & 0 \end{bmatrix} g_2^t \right) \xrightarrow{(\phi_1^-, \phi_2^-)} \left( \frac{g_{12}}{g_{14}}, \frac{g_{22}}{g_{24}} \right)$$

or

$$\left( \begin{bmatrix} 0 & 0 \\ g_1 & 1 \end{bmatrix} g_2^{-1}, \begin{bmatrix} 0 & 1 \\ g_1 & 0 \end{bmatrix} g_2^{-1} \right) \xrightarrow{(\phi_1^-, \phi_2^-)} \left( \frac{g_{12}}{g_{14}}, -\frac{g_{23}}{g_{21}} \right).$$

Let  $\varphi : M \longrightarrow \mathbb{H}_1^3(-1)$  be a timelike surface from an oriented and simply-connected open 2-manifold  $M$  into  $\mathbb{H}_1^3(-1)$  with  $ds_\varphi^2 = e^\omega(-dx^2 + dy^2) = e^\omega dudv$ . Then there exists an adapted framing  $F : M \longrightarrow \mathrm{SL}_2\mathbb{R} \times \mathrm{SL}_2\mathbb{R}$  of  $\varphi$

such that  $e_1 \circ F = e^{-\omega} \varphi_x$  and  $e_2 \circ F = e^{-\omega} \varphi_y$ . The generalized Gauß map  $\mathcal{G}$  of  $\varphi$  can be written

$$\begin{aligned} \mathcal{G} &= [(e_1 \circ F) \wedge (e_2 \circ F)] \\ &= [(e_1 + e_2)(F) \wedge (-e_1 + e_2)(F)] \\ &\cong \left( \left[ F_1 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} F_2^t \right], \left[ F_1 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} F_2^t \right] \right) : M \longrightarrow G(2, \mathbb{E}_2^4). \end{aligned}$$

Let  $\mathcal{G}_1 = \phi_1 \circ \mathcal{G}$  and  $\mathcal{G}_2 = \phi_2 \circ \mathcal{G}$ . Then  $\mathcal{G}_1 = \frac{F_{11}}{F_{13}}$ ,  $\mathcal{G}_2 = \frac{F_{21}}{F_{23}}$ . Note that the ordered pair  $(\mathcal{G}_1, \mathcal{G}_2) = \left( \frac{F_{11}}{F_{13}}, \frac{F_{21}}{F_{23}} \right)$  is the same as the hyperbolic Gauß map

$$[(e_0 + e_3)(F)] = [F_1(\mathbf{1} + \mathbf{k}') F_2^t].$$

If  $\mathcal{G} \cong \left( \left[ F_1 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} F_2^{-1} \right], \left[ F_1 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} F_2^{-1} \right] \right)$ , then

$$(\mathcal{G}_1, \mathcal{G}_2) = \left( \frac{F_{11}}{F_{13}}, -\frac{F_{24}}{F_{22}} \right) = [F_1(\mathbf{1} + \mathbf{k}') F_2^{-1}] = [(e_0 + e_3)(F)].$$

Let us define  $\mathcal{G}^- : M \longrightarrow G(2, \mathbb{E}_2^4)^-$  by

$$\begin{aligned} \mathcal{G}^- &= [(-e_1 + e_2)(F) \wedge (e_1 + e_2)(F)] \\ &\cong \left( \left[ F_1 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} F_2^t \right], \left[ F_1 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} F_2^t \right] \right). \end{aligned}$$

Let  $\mathcal{G}_1^- := \phi_1^- \circ \mathcal{G}^-$ ,  $\mathcal{G}_2^- := \phi_2^- \circ \mathcal{G}^-$ . Then

$$(\mathcal{G}_1^-, \mathcal{G}_2^-) = \left( \frac{F_{12}}{F_{14}}, \frac{F_{22}}{F_{24}} \right) = [F_1(\mathbf{1} - \mathbf{k}') F_2^t] = [(e_0 - e_3)(F)].$$

If  $\mathcal{G}^- \cong \left( \left[ F_1 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} F_2^{-1} \right], \left[ F_1 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} F_2^{-1} \right] \right)$ , then

$$(\mathcal{G}_1^-, \mathcal{G}_2^-) = \left( \frac{F_{12}}{F_{14}}, -\frac{F_{23}}{F_{21}} \right) = [F_1(\mathbf{1} - \mathbf{k}') F_2^{-1}] = [(e_0 - e_3)(F)].$$

### 13. THE LORENTZ HOLOMORPHICITY OF HYPERBOLIC GAUSS MAP AND TIMELIKE $\text{cmc} \pm 1$ SURFACES IN $\mathbb{H}_1^3(-1)$

In this section, we study the relationship between Lorentz holomorphicity of (projected) hyperbolic Gauß map and timelike  $\text{cmc} \pm 1$  surfaces in  $\mathbb{H}_1^3(-1)$ . Their relationship is summarized as the following theorem. Here, we assume that  $\varphi := \Phi_1 \Phi_2^t : M \longrightarrow \mathbb{H}_1^3(-1)$  and  $\psi := \Psi_1 \Psi_2^{-1} : M \longrightarrow \mathbb{H}_1^3(-1)$  are timelike surfaces in  $\mathbb{H}_1^3(-1)$ , where  $\Phi := (\Phi_1, \Phi_2) : M \longrightarrow \text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  and  $\Psi := (\Psi_1, \Psi_2) : M \longrightarrow \text{SL}_2\mathbb{R} \times \text{SL}_2\mathbb{R}$  are solutions of Lax equations (45) and (46), resp., in Theorem 4. Let us regard  $[(e_0 + e_3)(\Phi)] = [\Phi_1(\mathbf{1} + \mathbf{k}') \Phi_2^t]$   $[(e_0 + e_3)(\Psi)] = [\Psi_1(\mathbf{1} + \mathbf{k}') \Psi_2^{-1}]$  as the projected hyperbolic Gauß map

$\left(\frac{\Phi_{11}}{\Phi_{13}}, \frac{\Phi_{21}}{\Phi_{23}}\right) \in \mathbb{E}_1^2(u, v)$  ( $\left(\frac{\Psi_{11}}{\Psi_{13}}, -\frac{\Psi_{24}}{\Psi_{22}}\right) \in \mathbb{E}_1^2(u, v)$ ). Also, regard  $[(e_0 - e_3)(\Phi)] = [\Phi_1(1 - \mathbf{k}')\Phi_2']$  ( $[(e_0 - e_3)(\Psi)] = [\Psi_1(1 - \mathbf{k}')\Psi_2^{-1}]$ ) as the projected hyperbolic Gauß map  $\left(\frac{\Phi_{12}}{\Phi_{14}}, \frac{\Phi_{22}}{\Phi_{24}}\right) \in \mathbb{E}_1^2(u, v)$  ( $\left(\frac{\Psi_{12}}{\Psi_{14}}, -\frac{\Psi_{23}}{\Psi_{21}}\right) \in \mathbb{E}_1^2(u, v)$ ). Then we have the following theorem holds.

- Theorem 9.** (1)  $[(e_0 + e_3)(\Phi)]$  ( $[(e_0 + e_3)(\Psi)]$ ) is Lorentz antiholomorphic if and only if  $\varphi(\psi)$  satisfies  $H = 1$  and  $Q = 0$ .  
 (2)  $[(e_0 + e_3)(\Phi)]$  ( $[(e_0 + e_3)(\Psi)]$ ) is Lorentz holomorphic if and only if  $\varphi(\psi)$  satisfies  $H = 1$  and  $R = 0$ .  
 (3)  $[(e_0 - e_3)(\Phi)]$  ( $[(e_0 - e_3)(\Psi)]$ ) is Lorentz antiholomorphic if and only if  $\varphi(\psi)$  satisfies  $H = -1$  and  $Q = 0$ .  
 (4)  $[(e_0 - e_3)(\Phi)]$  ( $[(e_0 - e_3)(\Psi)]$ ) is Lorentz holomorphic if and only if  $\varphi(\psi)$  satisfies  $H = -1$  and  $R = 0$ .

*Proof.* We prove only part (1). The rest can be proved similarly. Since

$$[(e_0 + e_3)(\Phi)] = \left(\frac{\Phi_{11}}{\Phi_{13}}, \frac{\Phi_{21}}{\Phi_{23}}\right),$$

$[(e_0 + e_3)(\Phi)]$  is Lorentz antiholomorphic, i.e.,  $[(e_0 + e_3)(\Phi)]_u = 0$  if and only if  $(\Phi_{11})_u \Phi_{13} - \Phi_{11}(\Phi_{13})_u = 0$  and  $(\Phi_{21})_u \Phi_{23} - \Phi_{21}(\Phi_{23})_u = 0$ .

On the other hand, from the Lax equations (45),

$$\begin{aligned} \begin{pmatrix} (\Phi_{11})_u & (\Phi_{12})_u \\ (\Phi_{13})_u & (\Phi_{14})_u \end{pmatrix} &= \begin{pmatrix} \Phi_{11} & \Phi_{12} \\ \Phi_{13} & \Phi_{14} \end{pmatrix} \begin{pmatrix} \frac{\omega_u}{4} & \frac{1}{2}e^{\frac{\omega}{2}}(H+1) \\ -e^{-\frac{\omega}{2}}Q & -\frac{\omega_u}{4} \end{pmatrix} \\ &= \begin{pmatrix} \frac{1}{4}\Phi_{11}\omega_u - \Phi_{12}e^{-\frac{\omega}{2}}Q & \frac{1}{2}\Phi_{11}e^{\frac{\omega}{2}}(H+1) - \frac{1}{4}\Phi_{12}\omega_u \\ \frac{1}{4}\Phi_{13}\omega_u - \Phi_{14}e^{-\frac{\omega}{2}}Q & \frac{1}{2}\Phi_{13}e^{\frac{\omega}{2}}(H+1) - \frac{1}{4}\Phi_{14}\omega_u \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} \begin{pmatrix} (\Phi_{21})_u & (\Phi_{22})_u \\ (\Phi_{23})_u & (\Phi_{24})_u \end{pmatrix} &= \begin{pmatrix} \Phi_{21} & \Phi_{22} \\ \Phi_{23} & \Phi_{24} \end{pmatrix} \begin{pmatrix} -\frac{\omega_u}{4} & e^{-\frac{\omega}{2}}Q \\ -\frac{1}{2}e^{\frac{\omega}{2}}(H-1) & \frac{\omega_u}{4} \end{pmatrix} \\ &= \begin{pmatrix} -\frac{1}{4}\Phi_{21}\omega_u - \frac{1}{2}\Phi_{22}e^{\frac{\omega}{2}}(H-1) & \Phi_{21}e^{-\frac{\omega}{2}}Q + \frac{1}{4}\Phi_{22}\omega_u \\ -\frac{1}{4}\Phi_{23}\omega_u - \frac{1}{2}\Phi_{24}e^{\frac{\omega}{2}}(H-1) & \Phi_{23}e^{-\frac{\omega}{2}}Q + \frac{1}{4}\Phi_{24}\omega_u \end{pmatrix}. \end{aligned}$$

Thus,

$$\begin{aligned} (\Phi_{11})_u \Phi_{13} - \Phi_{11}(\Phi_{13})_u &= e^{-\frac{\omega}{2}}Q, \\ (\Phi_{21})_u \Phi_{23} - \Phi_{21}(\Phi_{23})_u &= \frac{1}{2}e^{\frac{\omega}{2}}(H-1). \end{aligned}$$

It then follows immediately that  $[(e_0 + e_3)(\Phi)]$  is Lorentz antiholomorphic if and only if  $\varphi$  satisfies  $H = 1$  and  $Q = 0$ .  $\square$

**Corollary 10.** (1)  $[(e_0 + e_3)(\Phi)]$  ( $[(e_0 + e_3)(\Psi)]$ ) is constant if and only if  $\varphi(\psi)$  satisfies  $H = 1$  and is totally umbilic ( $Q = 0$ , i.e.,  $Q = R = 0$ ).

- (2)  $[(e_0 - e_3)(\Phi)]$   $[(e_0 - e_3)(\Psi)]$  is constant if and only if  $\varphi$  ( $\psi$ ) satisfies  $H = -1$  and is totally umbilic.

#### 14. APPENDIX I: THE LAWSON-GUICHARD CORRESPONDENCE BETWEEN TIMELIKE cmc SURFACES IN DIFFERENT SEMI-RIEMANNIAN SPACE FORMS

In section 5, we discussed the *Lawson-Guichard correspondence* or simply *Lawson correspondence* between timelike cmc surfaces in semi-Riemannian space forms  $\mathbb{E}_1^3$ ,  $\mathbb{S}_1^3(1)$  and  $\mathbb{H}_1^3(-1)$ . In fact, this Lawson correspondence was already known to A. Fujioka and J. Inoguchi ([8] and [9]). In this appendix, we study the Lawson correspondence in a more general setting.

Let  $\bar{M}$  be a semi-Riemannian manifold and  $M \subset \bar{M}$  a hypersurface with the sectional curvatures  $\bar{K}$  and  $K$ , resp. Let  $\mathcal{S}$  be the shape operator derived from the unit normal vector field  $N$  on the hypersurface  $M$ . If  $X, Y$  span a nondegenerate tangent plane on  $M$ , then the Gauß equation is given by

$$(66) \quad K(X, Y) = \bar{K}(X, Y) + \epsilon \frac{\langle \mathcal{S}(X), X \rangle \langle \mathcal{S}(Y), Y \rangle - \langle \mathcal{S}(X), Y \rangle^2}{\langle X, X \rangle \langle Y, Y \rangle - \langle X, Y \rangle^2},$$

where  $\epsilon = \langle N, N \rangle$ . (See B. O'Neill [20] on p. 107.) We begin with the following theorem which can be found, for example, in T. Weinstein [24] on p. 158.

**Theorem 11** (Fundamental Theorem of Surface Theory: Lorentzian Version). *Given a simply-connected Lorentz surface with global null coordinate system, there exists a timelike immersion with the first and the second fundamental forms  $I$  and  $II$  if and only if  $I$  and  $II$  satisfy the Gauß and Mainardi-Codazzi equations.*

Let  $\mathcal{M}^3(\bar{K})$  be the semi-Riemannian 3-manifold with constant sectional curvature  $\bar{K}$ . For example,  $\mathcal{M}^3(-1) = \mathbb{H}_1^3(-1)$ ,  $\mathcal{M}^3(0) = \mathbb{E}_1^3$ , and  $\mathcal{M}^3(1) = \mathbb{S}_1^3(1)$ . For a conformal timelike immersion  $\varphi : M \rightarrow \mathcal{M}^3(\bar{K})$  with induce metric  $\langle \cdot, \cdot \rangle$ , Levi-Civita connection  $\nabla$ , Gaußian curvature  $K$  and shape operator  $\mathcal{S}$ , the Gauß and Mainardi-Codazzi<sup>9</sup> equations are satisfied:

$$(67) \quad K - \bar{K} = \det \mathcal{S} \text{ (Gauß Equation)}$$

$$(68) \quad (\nabla_X \mathcal{S})(Y) = (\nabla_Y \mathcal{S})(X) \text{ (Mainardi-Codazzi Equation)}$$

for all smooth vector fields  $X, Y, Z \in T(M)$ .

Assume that  $\varphi$  has a constant mean curvature  $H = \frac{1}{2} \text{tr}(\mathcal{S})$ . For any  $c \in \mathbb{R}$ , define

$$\tilde{\mathcal{S}} = \mathcal{S} + c\mathcal{I}, \quad \tilde{K} = \bar{K} - c \text{tr}(\mathcal{S}) - c^2,$$

---

<sup>9</sup>Since  $\mathcal{M}^3(\bar{K})$  has a constant sectional curvature and  $M$  is a hypersurface immersed into  $\mathcal{M}^3(\bar{K})$ , the Mainardi-Codazzi equation becomes (68). See B. O'Neill [20] on p. 115 for more details.

where  $\mathcal{I}$  is the identity transformation. Then the Gauß equation (67) and the Mainardi-Codazzi equation (68) still hold when  $\mathcal{S}$  and  $\bar{K}$  are replaced by  $\tilde{\mathcal{S}}$  and  $\tilde{K}$ , resp.:

$$\begin{aligned} K - \tilde{K} &= K - (\bar{K} - c \operatorname{tr}(\mathcal{S}) - c^2) \\ &= K - \bar{K} + c \operatorname{tr}(\mathcal{S}) + c^2 \\ &= \det(\mathcal{S}) + c \operatorname{tr}(\mathcal{S}) + c^2 \\ &= \det(\mathcal{S} + c\mathcal{I}) \\ &= \det(\tilde{\mathcal{S}}). \end{aligned}$$

Note that the Gaussian curvature  $K$  of  $M$  is intrinsic and does not change.

$$\begin{aligned} (\nabla_X \mathcal{S})(Y) &= \nabla_X \mathcal{S}(Y) - \mathcal{S}(\nabla_X Y), \\ (\nabla_Y \mathcal{S})(X) &= \nabla_Y \mathcal{S}(X) - \mathcal{S}(\nabla_Y X). \end{aligned}$$

Since  $(\nabla_X \mathcal{S})(Y) = (\nabla_Y \mathcal{S})(X)$ ,

$$\begin{aligned} \mathcal{S}([X, Y]) &= \mathcal{S}(\nabla_X Y - \nabla_Y X) \\ &= \mathcal{S}(\nabla_X Y) - \mathcal{S}(\nabla_Y X) \\ &= \nabla_X \mathcal{S}(Y) - \nabla_Y \mathcal{S}(X). \end{aligned}$$

Now,

$$\begin{aligned} \tilde{\mathcal{S}} &= \mathcal{S}([X, Y]) + c[X, Y] \\ &= \nabla_X \mathcal{S}(Y) - \nabla_Y \mathcal{S}(X) + c[X, Y] \\ &= \nabla_X (\mathcal{S} + c\mathcal{I})(Y) - \nabla_Y (\mathcal{S} + c\mathcal{I})(X) \\ &= \nabla_X \tilde{\mathcal{S}}(Y) - \nabla_Y \tilde{\mathcal{S}}(X). \end{aligned}$$

Therefore, there exists an immersion  $\tilde{\varphi} : M \longrightarrow \mathcal{M}^3(\tilde{K})$  with induced metric  $\langle \cdot, \cdot \rangle$  and shape operator  $\tilde{\mathcal{S}}$ , and  $\tilde{\varphi}(M)$  is isometric to  $\varphi(M)$ . The mean curvature  $\tilde{H}$  of  $\tilde{\varphi}(M)$  is

$$\tilde{H} = \frac{1}{2} \operatorname{tr}(\tilde{\mathcal{S}}) = \frac{1}{2} \operatorname{tr}(\mathcal{S}) + c = H + c$$

and this shows the Lawson-Guichard correspondence between timelike cmc  $H$  surfaces in  $\mathcal{M}^3(\bar{K})$  and timelike cmc  $(H + c)$  surfaces in  $\mathcal{M}^3(\tilde{K}) = \mathcal{M}^3(\bar{K} - 2cH - c^2)$ . In particular, when  $H = \bar{K} = 0$  and  $c = 1$ , we have the Lawson-Guichard correspondence between timelike minimal surfaces in  $\mathbb{E}_1^3$  and timelike cmc 1 surfaces in  $\mathbb{H}_1^3(-1)$ . If  $H = 0$  and  $\bar{K} = c = 1$ , then we have the Lawson-Guichard correspondence between timelike minimal surfaces in de Sitter 3-space  $\mathbb{S}_1^3(1)$  and timelike cmc 1 surfaces in  $\mathbb{E}_1^3$ .

15. APPENDIX II: SOME EXAMPLES OF TIMELIKE  $\text{cmc } \pm 1$  SURFACES IN  $\mathbb{H}_1^3(-1)$

In this appendix, we present some examples of timelike  $\text{cmc } 1$  surfaces in  $\mathbb{H}_1^3(-1)$ .

Let us consider the following stereographic projections in order to view the isometric images of timelike  $\text{cmc } \pm 1$  surfaces in  $\mathbb{H}_1^3(-1)$  into the interior  $\text{Int}\mathbb{S}_1^2(1) = \{(x_1, x_2, x_3) \in \mathbb{E}_1^3 : -(x_1)^2 + (x_2)^2 + (x_3)^2 < 1\}$  of de Sitter 2-space  $\mathbb{S}_1^2(1)$ .

Let  $\wp_+ : \mathbb{H}_1^3(-1) \setminus \{x_0 = -1\} \longrightarrow \mathbb{E}_1^3 \setminus \mathbb{S}_1^2(1)$  be the stereographic projection from  $-e_0 = (-1, 0, 0, 0)$ . Then

$$(69) \quad \wp_+(x_0, x_1, x_2, x_3) = \left( \frac{x_1}{1+x_0}, \frac{x_2}{1+x_0}, \frac{x_3}{1+x_3} \right).$$

Let  $\wp_- : \mathbb{H}_1^3(-1) \setminus \{x_0 = 1\} \longrightarrow \mathbb{E}_1^3 \setminus \mathbb{S}_1^2(1)$  be the stereographic projection from  $e_0 = (1, 0, 0, 0)$ . Then

$$(70) \quad \wp_-(x_0, x_1, x_2, x_3) = \left( \frac{x_1}{1-x_0}, \frac{x_2}{1-x_0}, \frac{x_3}{1-x_3} \right).$$

Cut  $\mathbb{H}_1^3(-1)$  into two halves by the hyperplane  $x_0 = 0$ . Denote by  $\mathbb{H}_1^3(-1)_+$  ( $\mathbb{H}_1^3(-1)_-$ ) the half containing  $e_0 = (1, 0, 0, 0)$  ( $-e_0 = (-1, 0, 0, 0)$ ). Then  $\wp_+ : \mathbb{H}_1^3(-1)_+ \longrightarrow \text{Int}\mathbb{S}_1^2(1)$  and  $\wp_- : \mathbb{H}_1^3(-1)_- \longrightarrow \text{Int}\mathbb{S}_1^2(1)$ .

*Example 1* (Timelike Enneper Cousin in  $\mathbb{H}_1^3(-1)$  (Isothermic Type)). Let  $(q, r) = (u, v)$ . Then using the Bryant-Umehara-Yamada type representation (62), we set up the following initial value problem:

$$F_1^{-1}dF_1 = \begin{pmatrix} u & -u^2 \\ 1 & -u \end{pmatrix} du, \quad F_2^{-1}dF_2 = \begin{pmatrix} v & -v^2 \\ 1 & -v \end{pmatrix} dv$$

with the initial condition  $F_1(0) = F_2(0) = \mathbf{1}$ . This initial value problem has a unique solution

$$F_1(u, v) = \begin{pmatrix} \cosh u & \sinh u - u \cosh u \\ \sinh u & \cosh u - u \sinh u \end{pmatrix},$$

$$F_2(u, v) = \begin{pmatrix} \cosh v & \sinh v - v \cosh v \\ \sinh v & \cosh v - v \sinh v \end{pmatrix}$$

which are Lorentz holomorphic and Lorentz antiholomorphic null immersions into  $\text{SL}_2\mathbb{R}$ . The Bryant type representation formula (51) then yields a time-like  $\text{cmc } 1$  surface in  $\mathbb{H}_1^3(-1)$ . The resulting surface is a correspondent of isothermic type<sup>10</sup> timelike Enneper surface in  $\mathbb{E}_1^3$  under the Lawson-Guichard correspondence. For this reason, the resulting surface is called *isothermic type timelike Enneper cousin* in  $\mathbb{H}_1^3(-1)$ .

---

<sup>10</sup>For details about isothermic and anti-isothermic timelike surfaces, please see [9] or [13].

Figure 1 shows different views of isothermic type timelike Enneper cousin in  $\mathbb{H}_1^3(-1)$  projected via  $\wp_+$  into the interior of the boundary  $\mathbb{S}_1^2(1)$ .




FIGURE 1. Isothermic type timelike Enneper cousin projected into  $\text{Int}\mathbb{S}_1^2(1)$  via  $\wp_+$  with light cone and the boundary  $\mathbb{S}_1^2(1)$  in  $\mathbb{E}_1^3$

Figure 2 shows different views of isothermic type timelike Enneper surface in  $\mathbb{E}_1^3$ .




FIGURE 2. Isothermic type timelike Enneper surface in  $\mathbb{E}_1^3$  with light cone

*Example 2* (Timelike Enneper Cousin in  $\mathbb{H}_1^3(-1)$  (Anti-isothermic Type)). Let  $(q, r) = (-u, v)$ . Then using the Bryant-Umehara-Yamada type representation (62), we set up the following initial value problem:

$$F_1^{-1}dF_1 = \begin{pmatrix} -u & -u^2 \\ 1 & u \end{pmatrix} du, \quad F_2^{-1}dF_2 = \begin{pmatrix} v & -v^2 \\ 1 & -v \end{pmatrix} dv$$

with the initial condition  $F_1(0) = F_2(0) = \mathbf{1}$ . This initial value problem has a unique solution

$$F_1(u, v) = \begin{pmatrix} \cos u & -\sin u + u \cos u \\ \sin u & \cos u + u \sin u \end{pmatrix},$$

$$F_2(u, v) = \begin{pmatrix} \cosh v & \sinh v - v \cosh v \\ \sinh v & \cosh v - v \sinh v \end{pmatrix}$$

which are Lorentz holomorphic and Lorentz antiholomorphic null immersions into  $\text{SL}_2\mathbb{R}$ . The Bryant type representation formula (51) then yields a timelike cmc 1 surface in  $\mathbb{H}_1^3(-1)$ . The resulting surface is a correspondent of anti-isothermic type timelike Enneper surface in  $\mathbb{E}_1^3$  under the Lawson-Guichard correspondence. For this reason, the resulting surface is called *anti-isothermic type timelike Enneper cousin* in  $\mathbb{H}_1^3(-1)$ .

Figure 3 shows different views of anti-isothermic type timelike Enneper cousin in  $\mathbb{H}_1^3(-1)$  projected via  $\wp_+$  into the interior of the boundary  $\mathbb{S}_1^2(1)$ .

Figures 4 and 5 show different views of anti-isothermic type timelike Enneper surface in  $\mathbb{E}_1^3$ .

*Example 3* (cmc 1 B-scroll in  $\mathbb{H}_1^3(-1)$ ). Let  $(q, r) = (u, 0)$ . Then using the Bryant-Umehara-Yamada type representation (62), we set up the following initial value problem:

$$F_1^{-1}dF_1 = \begin{pmatrix} u & -u^2 \\ 1 & -u \end{pmatrix} du, \quad F_2^{-1}dF_2 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} dv$$

with the initial condition  $F_1(0) = F_2(0) = \mathbf{1}$ . This initial value problem has a unique solution

$$F_1(u, v) = \begin{pmatrix} \cosh u & \sinh u - u \cosh u \\ \sinh u & \cosh u - u \sinh u \end{pmatrix},$$

$$F_2(u, v) = \begin{pmatrix} 1 & 0 \\ v & 1 \end{pmatrix}$$

which are Lorentz holomorphic and Lorentz antiholomorphic null immersions into  $\text{SL}_2\mathbb{R}$ . The Bryant type representation formula (51) yields a timelike cmc 1 surface

$$\varphi = F_1 F_2^t = \begin{pmatrix} \cosh u & -(u-v) \cosh u - \sinh u \\ \sinh u & -(u-v) \sinh u + \cosh u \end{pmatrix}$$

in  $\mathbb{H}_1^3(-1)$ . The resulting surface is a correspondent of minimal B-scroll in  $\mathbb{E}_1^3$  under the Lawson-Guichard correspondence.




FIGURE 3. Anti-isothermic type timelike Enneper cousin projected into  $\text{Int}\mathbb{S}_1^2(1)$  via  $\wp_+$  with light cone and the boundary  $\mathbb{S}_1^2(1)$  in  $\mathbb{E}_1^3$

Figure 6 shows different views of cmc 1 B-scroll projected in  $\mathbb{H}_1^3(-1)$  via  $\wp_+$  into the interior of the boundary  $\mathbb{S}_1^2(1)$ .

Figure 7 shows different views of minimal B-scroll in  $\mathbb{E}_1^3$ .




FIGURE 4. Anti-isothermic type timelike Enneper surface in  $\mathbb{E}_1^3$  with light cone

**Acknowledgment:** The author wishes to thank Dr. Jun-ichi Inoguchi for drawing his attention to this subject and for many invaluable suggestions. The author is grateful to Dr. Magdalena Toda for her hospitality during his short visit to Texas Tech University in 2003. Most of this paper grew out




FIGURE 5. Anti-isothermic type timelike Enneper surface in  $\mathbb{E}_1^3$  with self-intersection

of discussions with Dr. Jun-ichi Inoguchi who was also visiting Texas Tech University during that time.




FIGURE 6. cmc 1 B-scroll projected into  $\text{Int}\mathbb{S}_1^2(1)$  via  $\wp_+$  with light cone and the boundary  $\mathbb{S}_1^2(1)$  in  $\mathbb{E}_1^3$

# REFERENCES

- [1] R. Aiyama and K. Akutagawa, *Kenmostu-Bryant type representation formula for constant mean curvature spacelike surfaces in  $H_1^3(-c^2)$* , Differential Geom. Appl. **9** (1998), no. 3, 251–272.




FIGURE 7. Minimal B-scroll in  $\mathbb{E}_1^3$  with light cone

- [2] R. Aiyama and K. Akutagawa, *Kenmotsu-Bryant type representation formulas for constant mean curvature surfaces in  $H^3(-c^2)$  and  $S_1^3(c^2)$* , Ann. Global Anal. Geom. **17** (1999), no. 1, 49–75.
- [3] K. Akutagawa, *On spacelike hypersurfaces with constant mean curvature in the de Sitter space*, Math. Z. **196** (1987), 13–19.

- [4] R. L. Bryant, *Surfaces of mean curvature one in hyperbolic space*, Astérisque **154-155**, Soc. Math. de France, (1987), 321–347.
- [5] M. Dajczer and K. Nomizu, *On flat surfaces in  $S_1^3$  and  $H_1^3$* , Manifolds and Lie Groups (Notre Dame, Ind., 1980), 71–108, Progr. Math., 14, Birkhäuser, Boston, Mass., 1981.
- [6] J. Dorfmeister, J. Inoguchi and M. Toda, *Weierstraß-type representation of timelike surfaces with constant mean curvature*, Differential geometry and integrable systems (Tokyo, 2000), Contemp. Math., **308** (2002), Amer. Math. Soc., Providence, RI, 77–99.
- [7] C. Epstein, *The hyperbolic Gauss map and quasiconformal reflections*, J. Reine Angew. Math., **372** (1986), 96–135.
- [8] A. Fujioka and J. Inoguchi, *Timelike Bonnet surfaces in Lorentzian space forms*, Differential Geom. Appl. **18** (2003), no. 1, 103–111.
- [9] A. Fujioka and J. Inoguchi, *Timelike surfaces with harmonic inverse mean curvature*, In: Surveys on Geometry and Integrable systems, Advanced Studies in Pure Mathematics, Math. Soc. Japan, to appear.
- [10] J. Q. Hong, *Timelike surfaces with mean curvature one in anti-de Sitter 3-space*, Kodai Math. J., **17**, no.2, 1994, 341–350.
- [11] J. Inoguchi, *Timelike surfaces of constant mean curvature in Minkowski 3-space*, Tokyo J. Math., **21** (1998), no. 1, 141–152.
- [12] J. Inoguchi, *Private Communication*, 2003.
- [13] J. Inoguchi and M. Toda, *Timelike minimal surfaces via loop groups*, Acta. Appl. Math. **83** (2004), 313–335.
- [14] O. Kobayashi, *Maximal surfaces in the 3-dimensional Minkowski space  $\mathbb{L}^3$* , Tokyo J. Math. **6** (1983), no. 2, 297–309.
- [15] B. Lawson, *Complete minimal surfaces in  $S^3$* , Ann. Math. **92** (1970), 335–374.
- [16] S. Lee, *Spacelike surfaces of constant mean curvature one in de Sitter 3-space  $S_1^3$* , Illinois J. Math. **49**, Issue 1 (Spring 2005), 63–98.
- [17] M. A. Magid, *Timelike surfaces in Lorentz 3-space with prescribed mean curvature and Gauß map*, Hokkaido Math. J. **20** (1991), 447–464.
- [18] L. McNertney, *One-parameter families of surfaces with constant curvature in Lorentz 3-space*, Ph.D. Thesis, Brown Univ., Providence, RI, U.S.A., 1980.
- [19] T. K. Milnor, *A conformal analog of Bernstein's theorem in Minkowski 3-space*, The legacy of Sonya Kovalevskaya, Contemp. Math. **64** (1987), Amer. Math. Soc., Providence, RI, 123–130.
- [20] B. O'Neill, *Semi-Riemannian Geometry with Applications to Relativity*, Pure and Applied Mathematics, vol. 130, Academic Press, 1983.
- [21] B. Plamer, *Spacelike constant mean curvature surfaces in pseudo-Riemannian space forms*, Ann. Global Anal. Geom. **8** (1990), 217–226.
- [22] J. Ramanathan, *Complete spacelike hypersurfaces of constant mean curvature in de Sitter space*, Indiana Univ. Math. J. **36** (1987), 349–359.
- [23] M. Umehara and K. Yamada, *Complete surfaces of constant mean curvature-1 in the hyperbolic 3-space*, Annals of Math. **137** (1993), 611–638.
- [24] T. Weinstein, *An Introduction to Lorentz Surfaces*, de Gruyter Exposition in Math. 22, Walter de Gruyter, Berlin, 1996.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN MISSISSIPPI, HATTIESBURG, MS 39406-5045, U.S.A.

*E-mail address:* sunglee@usm.edu